

ASSOCIATION OF DENTAL PLAQUE, POOR ORAL HYGIENE AND PERIODONTAL DISEASE WITH HELICOBACTER PYLORI INFECTION

*A Dissertation Submitted
in partial fulfillment of the requirements
for the degree of*

MASTER OF DENTAL SURGERY

BRANCH – II

PERIODONTOLOGY



**THE TAMILNADU DR. M.G.R. MEDICAL UNIVERSTIY
CHENNAI-600 032
2007 – 2010**

Certificate

This is to certify that **Dr.SHEEJA JACOB**, Post graduate student (2007 – 2010) in the Department of Periodontology, Tamilnadu Government Dental College and Hospital, Chennai – 600 003 has done this dissertation titled “**ASSOCIATION OF DENTAL PLAQUE, POOR ORAL HYGIENE AND PERIODONTAL DISEASE WITH HELICOBACTER PYLORI INFECTION**” under our direct guidance and supervision in partial fulfillment of the regulations laid down by **The Tamil Nadu Dr.M.G.R. Medical University**, Chennai -600 032 for **M.D.S., (Branch – II) Periodontics** degree examination.

Dr.M.G. Krishna Baba
Professor & HOD

Dr. M.G. Krishna Baba
Professor & Guide

Department of Periodontology
Tamilnadu Government Dental College and Hospital
Chennai – 600 003.



Dr.K.S.G.A. NASSER
PRINCIPAL

Tamilnadu Government Dental College and Hospital
Chennai – 600 003.

I am extremely grateful to **DR.K. MALATHI M.D.S.**, Professor and HOD, Department of Periodontology, and **DR. S. KALAIVANI M.D.S.**, Professor, Tamilnadu Government Dental College and Hospital, Chennai, for their expert guidance and moral support during the completion of this study. I express my respect & deep sense of gratitude to **DR. M. G. KRISHNA BABA, M.D.S.**, Professor, Department of Periodontology, Tamilnadu Government Dental College and Hospital, who had the courage to stand by his beliefs. Without his unstinted guidance, support and encouragement, this study would not have been possible. Words cannot express my gratitude for his quiet confidence in my ability to do the study, his willingness to help clear the stumbling blocks along the way and his tremendous patience till the end of the study. I consider myself privileged, to have studied, worked and completed my dissertation under them in the Department.

I am also grateful to **DR. P.K. SARASWATHI M.D.S., DR. K.H.PANISHANKAR M.D.S.** Former Professors and HODs for the early start they gave and for their continual concern for our welfare.

My sincere thanks to **DR. K.S.G.A. NASSER, M.D.S.**, Principal Tamilnadu Government Dental College and Hospital, Chennai – 600 003, for his kind permission and encouragement.

I am extremely grateful to **DR.A.MUTHUKUMARASAMY, M.D.S., DR. M. JEEVA REKHA, M.D.S., DR.P.KAVITHA, M.D.S.**, Asst professors, **DR.P.BHUVANESWARI, M.D.S.**, former Asst. Professor Department of Periodontology, Tamilnadu Government Dental College

and Hospital, Chennai – 600 003 for their valuable suggestions, constant encouragement and timely help rendered throughout this study.

I thank **DR. MOHAMMED ALI, M.D., D.M.**, Professor and HOD, Department of Medical Gastroenterology, Government General Hospital, Chennai – 600 003 for granting me permission to conduct this study in the Department. At this juncture I would like to thank **DR. P.GANESH, M.D. D.M, DR. PREM KUMAR, M.D. D.M., DR. CAROLYN SELVI, M.D. D.M., DR. PADMANABHAN, M.D .D.M.**, Additional Professors and all post graduates of the Department of Medical Gastroenterology, Government General Hospital, Chennai for their kind co-operation in obtaining biopsy samples despite their routine busy schedule.

I extend my sincere thanks to **DR. SUMATHY PhD** Director, Department of Microbiology, Madras Medical College and Hospital, Chennai for permitting me to avail the laboratory facilities in the Department.

I am particularly thankful to **DR. S.CHITRA DCh, M.D.** Madras Medical College and Hospital, Chennai for co-partnering with me in the Dissertation work and for preparing the reagents and helping me at every stage of the Dissertation.

I wish to express my sincere thanks to **DR.C.R.SRIPRIYA, M.D.**, Assistant Professor, Department of Microbiology, Madras Medical College and Hospital, Chennai who took time off her busy schedule and expertly helped us in doing the ELISA test.

I take this opportunity to express my sincere gratitude to **DR.D.JAYANTHI M.D.S.**, for her constant encouragement, guidance and support throughout the study.

I thank **DR.A.K. MATHAI, Ph.D.**, Data Manager, National Institute of Epidemiology (ICMR) for helping me with the statistics in the study.

I also thank **Thiru SYED HUSSAIN, B.SC.**, of Citi Dot Net for typing and processing the dissertation.

I take this opportunity to express my gratitude to my colleagues and well wishers for their valuable help and suggestions throughout this study.

A special mention of thanks to all my patients for their consent, co-operation and participation in this study.

All glory and honour to **THE LORD ALMIGHTY** who gives me the strength to persist against all odds, whose loving kindness and mercies endureth forever.

My heartfelt and deep gratitude to all my family members, my in-laws for helping me out with my children, my parents and sisters for their help, patience, love and prayers which have sustained me throughout this period.

I sincerely thank my husband and children for silently and patiently encouraging me, enduring with me, praying for me and being a pillar of moral support and strength throughout this period.

ABSTRACT

Aim:

Peptic ulcer, the most common stomach disease is now well accepted as an infectious disease; and the causative agent *H. pylori* must be treated with antibiotics. Recently attention has been focused on the importance of dental plaque in harbouring *H.pylori* and its role as a potential reservoir for gastric infection & reinfection. This study was undertaken to determine whether dental plaque harbours *H. pylori* and to determine the association between oral hygiene and periodontal disease status and *H.pylori* gastritis

Materials and Methods:

110 patients with dyspeptic symptoms and clinical indications for an upper gastroendoscopy from the Department of Medical Gastroenterology, Government General Hospital Chennai were selected for the study. Oral hygiene status and periodontal disease status were examined and study variables were obtained from patient's history. Among the 110 patients, 55 patients who had a positive *H.pylori* serology or positive rapid urease test or histologic evidence for the presence of *H.pylori* in antral biopsy specimens were categorized as cases. The remaining 55 patients who were negative for these tests were controls. The presence of *H.pylori* in dental plaque was detected by the rapid urease test and culture.

Results:

It was found that the association of periodontal disease and poor oral hygiene with *H.pylori* gastritis was not significant. The RUT was positive in 87.3% of cases and 52.7% of controls. But there was no cultural detection of *H. pylori* in dental plaque.

Conclusion:

Hence the presence of *H.pylori* in dental plaque is inconclusive. Further studies using larger sample size and specific methods of detection of *H.pylori* are required to better assess the role of dental plaque as a reservoir for *H.pylori* and its relationship with *H.pylori* infection.

CONTENTS

S.NO.	TITLE	PAGE NO.
1.	INTRODUCTION	1
2.	AIM AND OBJECTIVES	3
3.	REVIEW OF LITERATURE	4
4.	MATERIALS AND METHODS	14
5.	RESULTS	43
6.	DISCUSSION	62
7.	SUMMARY AND CONCLUSION	70
8.	BIBLIOGRAPHY	72
9.	APPENDIX	81

LIST OF ABBREVIATIONS

1. AAP	–	American Academy of Periodontology
2. BOD	–	Biological oxygen demand
3. CAL	–	Clinical attachment level
4. CEJ	–	Cemento Enamel Junction
5. CI-S	–	Calculus Index- Simplified
6. CLO	–	Campylo-bacter like organism
7. DI-S	–	Debris Index- Simplified
8. ELISA	–	Enzyme linked immunosorbent assay
9. H. pylori	–	Helicobacter pylori
10. IgG	–	Immunoglobulin G
11. MALT	–	Mucosa associated lymphoid tissue
12. NHANES	–	National Health and Nutrition Examination Survey
13. NSAIDs	–	Non-steroidal anti-inflammatory drugs
14. OHI-S	–	Oral hygiene index- Simplified
15. PPD	–	Probing pocket depth
16. RU/ml	–	Relative units/ml
17. RUT	–	Rapid urease test
18. RUT(P)	–	Rapid urease test(plaque)
19. RUT(B)	–	Rapid urease test(biopsy)
20. SES	–	Socioeconomic status
21. SDS	–	Sodium dodecyl sulphate

LIST OF PHOTOGRAPHS

S. No.	Title	Page No
1.	Good oral hygiene in H.pylori positive patient	32
2.	Fair oral hygiene in H.pylori positive patient	32
3.	Poor oral hygiene in H.pylori positive patient	33
4.	Periodontitis in H.pylori positive patient	33
5.	Armamentarium for dental plaque and blood sampling	34
6.	Pocket depth > 5mm in H. pylori positive patient	34
7.	Aliquots with urea broth and saline for RUT and sample collection	35
8.	Transport for samples	35
9.	Collection of dental plaque samples	36
10.	Endoscope & biopsy forceps for antral biopsy	36
11.	Antral biopsy obtained during endoscopy by Gastroenterologist	37
12.	Antral biopsy taken from the biopsy forceps	37
13.	Collection of blood samples for ELISA	38
14.	Armamentarium for culture; streaking of sample onto culture plate	38
15.	Electron Microscope	39
16.	ELISA kit for Anti-H.pylori (Ig G)	39
17.	Serum samples for ELISA	40
18.	Microplate wells coated with antigen	40
19.	RUT positive in dental plaque and antral biopsy samples	59
20.	Culture positive for H.pylori in antral biopsy	59
21.	Giemsa stain showing H. pylori in antral biopsy specimen	60
22.	Histopathological features of chronic gastritis	60
23.	Microplate wells after ELISA reaction	61
24.	Microplate wells in ELISA reader	61

LIST OF TABLES

S. No.	Title	Page No
I	Master Chart-Cases	46
II	Master Chart- Controls	48
III	Comparison of gender between cases and controls	50
IV	Comparison of handling of animals between cases and controls	51
V	Comparison of Socioeconomic status between cases and controls	52
VI	Comparison of mean values between cases and controls	53
VII	Comparison of smoking habit and alcohol consumption between cases and controls	53
VIII	Comparison of OHI score between cases and controls	54
IX	Comparison of periodontal disease and pocket depth > 5mm between cases and controls	55
X	Comparison of RUT(P) and culture(P) between cases and controls	56
XI	Comparison of RUT(B), culture(B), Gram and Giemsa stain between cases and controls	57
XII	Results of multiple logistic regression analysis	57
XIII	Association of OHI and periodontal disease with RUT(P) in cases	58
XIV	Association of OHI and periodontal disease with RUT(P) in controls	58

LIST OF FIGURES

S. No.	TITLE	PAGE NO
I	Distribution of gender in cases and controls	50
II	Comparison of handling of animals between cases and controls	51
III	Comparison of Socioeconomic status between cases and controls	52
IV	Comparison of OHI score between cases and controls	54
V	Comparison of periodontal disease between cases and controls	55
VI	Comparison of RUT(P) between cases and controls	56

INTRODUCTION

The stomach contains about 10^4 bacterial cells/ml of gastric content (*Marianne Quiding – Jabrink et al 2009*).⁶⁷ The comparatively low number of bacteria in the stomach results from the low pH (pH-2) caused by the production of hydrochloric acid and a considerable flow (the turnover rate is 4-6 l/hr) (*Tannock 1995*).¹⁰⁴ Until recently, the stomach was considered to be without an established normal flora. *H.pylori*, a spiral shaped, motile gram – negative, micro – aerophilic bacterium was then discovered (*Marshall BJ, Warren JR 1984*).⁷⁰

It is now well accepted that the most common stomach disease, peptic ulcer is an infectious disease and that the causative agent, *H. pylori* must be treated with antibiotics (*European Study Group 1997*).³¹ Stomach infection with this organism causes inflammation of the gastric mucosa which can lead to gastritis, duodenal or gastric ulcers and even in rare cases to gastric carcinoma or MALT (mucosa associated lymphoid tissue) lymphoma (*Czenikiewicz- Guzik et al 2004*).²²

Helicobacter pylori is one of the most common bacterial infections in humans (*Blaser 1997*).¹⁵ Approximately 50% of the world's population is believed to be infected with *H.pylori* (*Czenikiewicz- Guzik et al 2004*).²² Serological tests have shown that the carriage rate of *H.pylori* is reported to be 20 – 80% for adults in the developed world and this figure may rise to more than 90% in the developing world (*Taylor & Blaser 1997*).¹⁰⁵ The majority of infected individuals do not develop clinically apparent disease, but there is now indisputable evidence that 6-20% of infection results in peptic ulceration and a smaller proportion (less than 1%) are associated with gastric cancer. Infection in a given individual will result either in the peptic ulcer pathway with associated increased acid output or the chronic atrophic

gastritis – carcinoma pathway which is associated with hypo or achlorhydria (*Michael G. Farthing 1998*).⁷⁹

Although eradication of *H.pylori* can be achieved with a combination therapy of antibiotics, the possibility of recurrence is very high (*Unge P 1996*).¹⁰⁷ The reservoir of *H. pylori* and its mode of transmission are unclear. A faecal- oral, oral-oral and gastro- oral route of infection have been suggested (*Mai et al 1989*).⁶¹ Recently researchers have suggested that the primary extra gastric reservoir for *H.pylori* is the oral cavity (*Nabwera, Logan 1999*).⁸⁴ *H.pylori* has been detected by various methods in dental plaque,^{49,62,25,7,52,65} which has led to the suggestion that dental plaque may be responsible for the transmission of the bacteria and possibly serve as a source of reinfection after eradication treatment. However some studies^{21, 22, 94} have reported no correlation between dental presentation of the micro organism and *H.pylori* associated gastritis. The hypothesis that oral flora may be a permanent reservoir of viable *H.pylori* is still inconclusive.

Since human infection by this pathogen appears to involve an oral route, it seems biologically plausible that oral health status directly or indirectly influences the process of *H.pylori* infection or reinfection.

This study was undertaken (a) to determine whether dental plaque harbors *H.pylori* and may act as a reservoir for the organism and (b) to determine the relation between *H.pylori* gastritis and dental health.

AIM AND OBJECTIVES

The aim of the present study was to determine whether dental plaque harbors *H. pylori* and thereby act as a reservoir for the organism and to determine the relation between oral hygiene status and periodontal disease status with *H. pylori* infection.

For this purpose the following objectives were undertaken:

1. To determine the presence of *H. pylori* in dental plaque of patients with dyspeptic symptoms and indications for gastroendoscopy
2. To determine the presence of *H. pylori* in dental plaque of patients with good, fair and poor oral hygiene and correlate with *H. pylori* in antral biopsy of the same patients.
3. To determine the presence of *H. pylori* in periodontitis patients and correlate with *H. pylori* infection.

REVIEW OF LITERATURE

The discovery of *Helicobacter pylori* in 1982 by **Warren and Marshall 1983**¹¹² was the starting point of a revolution concerning the concepts and management of gastro duodenal disease. The public health importance of *H.pylori* and its role in stomach diseases was recognized in 2005 by the attribution of the Nobel Prize in Physiology or Medicine to **B.Marshall** and **R.Warren**. In the history of Noble Prizes, this is only the third time that the discovery of a bacterium has been acknowledged.³⁵

PRE H. PYLORI ERA

Donati M. 1586²⁶ gave the first description of gastric ulceration.

Morgagn G.B. 1761⁸² described erosions and erythema of the stomach and duodenum in patients with heartburn and upper abdominal pain.

Billroth.CAT. 1881¹¹ described gastric surgeries- gastroduodenostomy and gastrojejunostomy.

Von Mikulicz – Radecki J. 1881¹¹¹ reported the first potentially usable endoscope.

Schwarz K. 1910¹⁰⁰ gave the famous dictum “no acid, no ulcer”

Sippy B.W. et al 1915¹⁰¹ recommended antacids in the treatment of peptic ulcer.

Ash. ASF. Schild H.O. 1966³ described the gastric histamine receptor.

Black J.W. et al 1972¹⁴ reported the first H₂ receptor antagonist.

*Ganser A.L., Forte J.G. 1973*³⁷ demonstrated the presence of a potassium stimulated adenosine triphosphate pump. Six years later the first commercially available proton pump inhibitor was reported.

KEY DATES IN HISTORY OF H. PYLORI

- 1893¹³ - Gastric spiral bacteria were reported for the first time in the stomach of dogs
- 1906⁵³ - Spirochetes were demonstrated in the human stomach
- 1924⁵⁸ - Urease activity in the stomach was reported
- 1950³³ - Urease in patients with gastric ulceration was reported to neutralize gastric acid via the production of ammonia.
- 1954⁹² - A study of 1000 gastric biopsy specimens failed to confirm the presence of bacteria
- 1975¹⁰³ - Gastric spirochetes and gastritis were present in 80% of gastric ulcers
- 1983¹¹² - **Discovery of H. Pylori.** Campylobacter – like organisms associated with gastritis and possibly peptic ulceration – beginning of modern era. This was first presented at the Royal Australian College of Physicians on 22nd October 1982 and published in letter form in 1983.
- 1985⁶⁹ - Temporal relationship was established between acquisition of H.pylori and development of gastritis.

- 1987²⁰ - It was proposed that eradication of H.pylori leads to long – term cure of duodenal ulceration
- 1989^{40,30} - The genus ‘Helicobacter’ was suggested.
- 1994⁴⁶ - H.pylori was classified as a grade I (definite carcinogen)
- 1994⁸⁸ - It was suggested that the infection should be eradicated in patients with peptic ulcers.
- 1997⁶⁴ - European consensus report was given on the management of H.pylori infection.

EPIDEMIOLOGY OF H.PYLORI

H.pylori has been reported world wide, but no common extra human reservoir of H.pylori has been detected. Although prevalence is decreasing, H.pylori infection remains one of the most common in man.

Roger A. Feldman et al 1988⁹⁹ after analysis of data of several studies reported that the major period of H.pylori acquisition is in childhood.

Mendall et al 1992⁷⁸ found that crowded living conditions are also associated with increased carriage rates of H.pylori.

The Eurogast study group 1993¹⁰⁶, Gasbarrini G. et al 1995³⁸ reported that the seroprevalence of H.pylori infection is often similar in males and females.

Fox JG. 1995³⁴ reported that cats are found with H.pylori in nature and could be a reservoir for human infection.

*Isabelle. M. Madinier et al 1997*⁴⁷ found that in developing countries, almost all children are infected by age 10. The major risk factors are poor sanitary conditions during childhood.

*Malaty.H.M. et al 1996*⁶³, *the Eurogast Study group 1993*¹⁰⁶, *Veldhuyzen van Santen S.J. 1995*¹¹⁰ in their studies found that lower socio – economic status and/or a low level of education are associated with an increase in prevalence of H.pylori infection.

*Roger A. Feldman et al 1998*⁹⁹ reported that consumption of alcohol associated with seropositivity, may be an indirect measure of socioeconomic and cultural variables.

*Dowsett S.A., M.J. Kowolik 2003*²⁷ reported that once H.pylori is acquired, it persists into old age, unless eradicated by treatment.

H.PYLORI IN DENTAL PLAQUE – ORAL CAVITY AS A RESERVOIR FOR H.PYLORI

The human stomach had long been considered a sterile organ unsuitable for microbial colonization because of its acidity

*Marshall FJ. et al 1984*⁷², *Marshall FJ et al 1987*⁷¹, *Goodwin CS et al 1989*⁴⁰ demonstrated that Helicobacter pylori was a gram – negative microaerophilic motile bacterium, especially adapted to life in the human stomach.

*Megraud F et al 1985*⁷⁷ found that H. pylori was capable of surviving and proliferating in this hostile environment by mobility in the gastric mucus (flagella, enzymatic mucus degradation), partial neutralization of gastric

acidity (urease converting urea into ammonia), and specific adherence to gastric epithelial cells.

Marshall et al 1985⁶⁹ reported that the mode of transmission of H.pylori remains something of a mystery. Since blood borne infection seems improbable, H.pylori must reach the stomach, its primary residence, via the oral cavity. **Marshall (1985)**,⁶⁹ in an attempt to convince the then many skeptics of the role of H.pylori in gastric disease, ingested H.pylori orally to confirm that it could indeed cause gastritis. He developed symptomatic gastritis thereby fulfilling Koch's postulates

Dowsett S.A., Kowolik. M.J. 2003²⁷ reported that to date, there has been little success in identifying a consistent non human source of infection, and it was likely that H.pylori is spread directly from person to person.

DETECTION OF H.PYLORI FROM ORAL SPECIMENS

Studies have shown frequent isolation of H.pylori from oral specimens particularly dental plaque.

H.pylori has been detected in subgingival plaque **Pustorino et al 1996**⁹⁵, **Dowsett et al 1999**²⁸, **Riggio and Lennon 1999**,⁹⁸ saliva and oral lesions **Mravak – Stipetic et al 1998**⁸³, **Birek et al 1999**¹² on oral mucosa **Mravak – Stipetic et al 1998**⁸³, **Dowsett et al 1999**²⁸, **Allaker et al 2002**,¹ and in supragingival plaque **Allaker et al 2002**¹, **Kim et al 2000**⁵¹, **Song et al 2000**.¹⁰²

Different methods have been used for the diagnosis of gastric infection and for the detection of oral H. pylori.

RAPID UREASE TEST

*Gill et al 1994*³⁶, *Pytko – Polonczyk et al 1996*⁹⁶, *Desai et al 1998*²⁵, *Butt et al 1999*¹⁷, and *Avcu et al 2001*⁵, *Pradeep et al 2006*⁹⁴ – These studies used the rapid urease test for the detection of oral H.pylori.

H.pylori is the only urease positive bacterium known to reside in the stomach; hence the rapid urease test provides a suitable detection method for use on gastric samples. The oral cavity is residence to several urease producing specimens including Streptococcus species, Haemophilus species and Actinomyces species. *Vaira et al 1998*¹⁰⁹ reported that the urease producing oral bacteria unlike H.pylori usually cannot give positive results within an hour.

CULTURE

*Krajden et al 1989*⁵² reported first the presence of H.pylori in oral cavity, when the bacterium was cultured from the dental plaque of one of 29 patients with H.pylori associated gastric disease. Since then culture of oral H.pylori has met with limited success.

The detection rates by culture in dental plaque samples have been consistently low. There are few exception studies.

*Majumdar et al 1990*⁶² had oral H.pylori culture positive samples in 40/40 patients and *D'Alessandro and Seri 1992*²³ isolated H.pylori in 16/ 20 patients by culture from dental plaque.

*Oshowo et al 1998*⁹⁰ isolated H.pylori by culture from dental plaque in 2 of 180 patients.

*Parsonnet et al 1999*⁹³ obtained 3 culture positive samples from a total of 26 patients.

*Namavar et al 2001*⁸⁵ – In both the studies with high culture positive results, confirmation of isolate identity by means of molecular methods was not performed and as a result there is a potential for false – positives.

*Allaker et al 2002*¹, *Luman et al 1996*⁵⁹, *Hardo et al 1995*⁴⁴ could not isolate H.pylori by culture from dental plaque - 0/109, 0/100, 0/62 patients respectively.

With the advances in molecular technology, the potential difficulties with culture have been circumvented by the use of the polymerase chain reaction *Dowsett et al 2003*.²⁷

PCR

PCR assays can detect the target DNA regardless of the viability of the bacteria *Mapstone NP et al 1993*⁶⁶.

*Yang H.T. 1993*¹¹³, *Mapstone NP et al 1993*⁶⁶, *Li.C. et al 1995*⁵⁷ demonstrated that nested PCR has a high sensitivity and specificity than one step PCR for samples with abundant bacterial flora

*Asikainen et al 1994*⁴, *Banatvala et al 1994*⁸ reported that different sets of primers are used based on the urease gene, 16S ribosomal RNA, genes encoding H.pylori species-specific proteins or randomly selected DNA fragments.

*Asikainen et al 1994*⁴ showed prevalence of 0% in dental plaque by PCR from a total of 336 samples, and *Riggio and Lennon 1999*⁹⁸ 38% of 29 samples in randomly selected patients, while *Banatvala et al 1993*⁷

showed a prevalence of 86% of 21 samples in symptomatic H.pylori positive patients.

ELISA

*Jones D.M., et al 1984*⁴⁹, *Marshall B.J 1984*⁷⁰ reported that the antibody response to H.pylori infection was used for the diagnosis of this infection immediately following the discovery of H.pylori.

*Mitchell et al 1987*⁸⁰ stated that H.pylori infection is a chronic condition and IgG (subclasses 1 and 4) is the predominant immunoglobulin class even in children. IgG are present at the mucosal level and detected in virtually all blood samples.

*Kupier.S.E. et al 1993*⁵⁴ reported that H.pylori almost constantly induces a specific systemic immune response which may reflect the antibodies produced at the gastric mucosal level, while only 2% of patients fail to seroconvert.

*Everhart JE et al 2000*³² found that the serological testing for H.pylori antibodies using an ELISA has a reported sensitivity and specificity of 91% and 97%.

RELATIONSHIP BETWEEN POOR ORAL HYGIENE, PERIODONTAL DISEASE ATTRIBUTES AND HELICOBACTER PYLORI INFECTION

*Nabwera HM, Logan RP 1999*⁸⁴ recently suggested that the primary extragastric reservoir for H.pylori is the oral cavity.

Since human infection by this pathogen appears to be involving an oral route, it seems biologically plausible that oral health status directly or indirectly influences the process of H.pylori infection or reinfection.

Bielanski 1999¹⁰ reported positive association between poor oral health and H.pylori. He performed a large epidemiological investigation using urea – breath test in over 10,000 subjects, which revealed that periodontitis significantly increases the risk of gastric H.pylori infection.

Riggio and Lennon 1999⁹⁸ reported in their study, that when subgingival plaque samples were obtained from periodontal pockets at least 5mm deep, 33% of the pocket sites analyzed were H.pylori positive and 38% of the 29 subjects with moderate to severe periodontal pocketing were positive for H.pylori.

Avcu .N. et al 2001⁵ correlated H.pylori positivity in dental plaque with OHI scores. The oral hygiene status was assessed by the Oral Hygiene Index of Greene and Vermillion. The positivity was 28.5%, 90.2% or 100% in patients with good, fair or poor OHI scores respectively.

Ozdemir A et al 2001⁹¹ evaluated the oral hygiene status of gastritis patients using the Quigley Hein plaque index. The study did not report on the relationship of plaque scores with H.pylori infection.

Berroteran A. et al 2002⁹ reported that there was no correlation between H.pylori infection and dental hygiene as assessed by the Plaque Index of Silness and Loe. He also reported no correlation between periodontal disease and H.pylori infection.

Dye.B A. et al 2002²⁹ reported from their study based on the data from the first phase of the Third National Health and Nutrition Examination survey, that periodontal disease characterized by deep periodontal pockets, may be associated with H.pylori infection in adults in the United States. They reported that periodontal pockets with a depth ≥ 5 mm were associated with increased odds of H pylori seropositivity after adjustment of sociodemographic factors.

Choudary CR et al 2003¹⁹ suggested that periodontal health would determine whether oral H.pylori may or may not cause gastric infection.

Pradeep S. Anand et al 2006⁹⁴ examined the oral hygiene status of patients using the Oral Hygiene Index of Greene and Vermillion. The oral hygiene status of the patients was classified into good, fair and poor depending on their oral hygiene scores. Among cases 34 had poor oral hygiene, 25 had fair and 6 had good oral hygiene. Among controls 28 had poor, 33 had fair and eight had good oral hygiene. The study reported little correlation between oral hygiene status and H.pylori infection.

In the above study the periodontal status of the patients was also examined as a dichotomous variable and the patients described as either healthy or diseased. Among cases 30 of 65 subjects had periodontal disease compared to only 20 of 69 subjects among controls. Although there were more periodontal disease subjects among cases than controls, the observed difference was not statistically significant and they concluded that there is no association between periodontal disease attributes and H. pylori infection.

MATERIALS AND METHODS

STUDY DESIGN AND SUBJECT SELECTION

On approval from the Institutional ethical committee, the study was conducted at the Department of Medical Gastroenterology, Govt. General Hospital, Chennai from November 2008 – June 2009. The study included 110 patients with various symptoms relating to the gastrointestinal tract most commonly epigastric pain and dyspepsia and clinical indications for an upper gastrointestinal endoscopy. The patients were of either sex and in the age range from 17 to 76 years.

A written informed consent was obtained from all patients. The details of complete history and clinical features of the subjects undergoing endoscopy were obtained.

Preprocedure preparations for oesophagogastroendoscopy were performed according to standard methods. Prior to endoscopy assessment of oral hygiene status, clinical parameters-probing depth and clinical attachment level were measured and plaque samples were obtained for identification of H.pylori by rapid urease test and culture.

Biopsies of gastric tissue were collected from the antrum of patients undergoing endoscopy and specimens were used for 1) histopathology study 2) culture 3) rapid urease test 4) Gram and Giemsa staining for identification of H.pylori infection.

Venous blood samples were collected from the patients for serological diagnosis of H.pylori infection.

INCLUSION CRITERIA

All patients with complaints suggestive of upper gastrointestinal disease –i.e.; dyspepsia and who were to undergo endoscopy for the same were included in the study.

EXCLUSION CRITERIA

1. Patients with active bleeding ulcers
2. Post gastrectomy individuals
3. Patients with history of proton pump inhibitor within 2 weeks of endoscopy or antibiotic intake within one month prior to the study.
4. Patients with history of chronic use of NSAIDS.
5. Patients who have undergone oral prophylaxis within past 6 months.
6. Patients on mouth rinses.

The subjects were categorized into two groups cases (n = 55) and controls (n = 55).

Subjects with clinical symptoms and a positive test for any of the three diagnostic tests (histopathology, or rapid urease test on antral biopsy specimens or serology) were cases.

Subjects with clinical symptoms and a negative test for H.pylori serology or negative rapid urease test and histopathology of antral biopsy specimen were controls.

STUDY PROTOCOL

- 1) Institutional Ethical committee approval

- 2) Medical history and informed consent
- 3) Assessment of oral hygiene status by Simplified Oral Hygiene Index –Greene and Vermilion 1964.⁴²
- 4) Periodontal examination using clinical parameters – probing depth and clinical attachment level.
- 5) Collection of plaque samples.
- 6) Collection of antral biopsy samples.
- 7) Collection of blood samples
- 8) Rapid urease test
- 9) Culture.
- 10) Histopathology
- 11) ELISA for detection of IgG antibody to H.pylori.
- 12) Statistical analysis

A detailed history and informed consent were obtained from all the subjects after explaining the study procedure.

The study variables obtained from the patient's history were age, gender, socioeconomic status, handling of animals, smoking(ex-smoker, current smoker, non-smoker) and alcohol consumption(current, past, never).

Socioeconomic status of the patient was classified as Upper, Upper middle, Lower middle; Upper lower and Lower based on Kuppuswamy's Socioeconomic Status Scale 2007. The original total scores for each of these groups is as given in the table.

KUPPUSWAMY'S SOCIOECONOMIC STATUS SCALE 2007⁵⁵

(A) Education	Score
----------------------	--------------

1. Profession or Honors	7
2. Graduate or post graduate	6
3. Intermediate or post high school diploma	5
4. High school certificate	4
5. Middle school certificate	3
6. Primary school certificate	2
7. Illiterate	1

(B) Occupation	Score
-----------------------	--------------

1. Profession	10
2. Semi-Profession	6
3. Clerical, Shop-owner, Farmer	5
4. Skilled worker	4
5. Semi-skilled worker	3
6. Unskilled worker	2
7. Unemployed	1

(C) Family income per Month (in Rs) - original	Score	Modified for 2007
1. =2000	12	=19575
2. 1000-1999	10	9788-19574
3. 750-999	6	7323- 9787
4. 500-749	4	4894- 7322
5. 300-499	3	2936-4893
6. 101-299	2	980-2935
7. =100	1	=979

TOTAL SCORE	SOCIOECONOMIC CLASS
26-29	Upper (I)
16-25	Upper Middle(II)
11-15 Middle	Lower middle (III)
5-10 Lower	Upper lower (IV)
<5	Lower(V)

In the present study the SES was classified as Upper, Middle and Lower because of small sample size. The total scores for these three groups that were used to arrive at the socioeconomic status of the patients were as follows: 26-29 Upper; 11-25 Middle; <5-10 Lower.

The oral hygiene status was assessed using the Simplified Oral Hygiene Index (OHI-S) Greene and Vermilion – 1964,⁴² using mouth mirror and explorer.

SIMPLIFIED ORAL HYGIENE INDEX (OHI – S) GREENE AND VERMILION 1964⁴²

Teeth examined – 16, 11, 26, 46, 31, and 36. If any of these teeth is missing, immediate distal tooth is examined.

Surfaces examined – buccal surfaces of upper molars, lingual surfaces of lower molars and labial surface of upper and lower incisors.

The OHI – S consists of 2 components.

A. Simplified Debris Index (DI – S)

B. Simplified Calculus Index (CI – S)

DEBRIS INDEX – SIMPLIFIED

SCORING CRITERIA

- | | |
|---|--|
| 0 | - No debris or stain present |
| 1 | - Soft debris covering $< 1/3^{\text{rd}}$ of the tooth surface or the presence of extrinsic stains without debris, regardless of the surface area covered |
| 2 | - Soft debris covering $> 1/3^{\text{rd}}$ but $< 2/3^{\text{rd}}$ of the exposed tooth surface. |
| 3 | - Soft debris covering $> 2/3^{\text{rd}}$ of the exposed tooth surface. |

$$\text{Debris index simplified score per person} = \frac{\text{Sum of debris score per tooth surface}}{\text{Number of surfaces examined}}$$

CALCULUS INDEX – SIMPLIFIED

SCORING CRITERIA

- 0 - No calculus present
- 1 - Supragingival calculus covering $< 1/3^{\text{rd}}$ of the exposed tooth surface.
- 2 - Supragingival calculus covering $> 1/3^{\text{rd}}$ but $< 2/3^{\text{rds}}$ of the exposed tooth surface (or)

The presence of individual flecks of sub gingival calculus around the cervical portion of the tooth (or) both.
- 3 - Supragingival calculus covering $> 2/3^{\text{rds}}$ of the exposed tooth surface (or)

A continuous heavy band of sub gingival calculus around the cervical portion of the tooth (or) both.

$$\text{Calculus index – simplified score per person} = \frac{\text{Sum of calculus score per tooth surface}}{\text{Number of teeth examined}}$$

$$\text{OHI – S score per person} = \text{DI - S} + \text{CI – S Score}$$

Interpretation

Good = 0.0 – 1.2

Fair = 1.3 – 3.0

Poor = 3.1 – 6.0

Periodontal evaluation was done by measuring probing depth (PPD) and clinical attachment level (CAL) using mouth mirror and Williams's periodontal probe. Based on AAP classification 1999,² patients with periodontal disease – chronic periodontitis – localized (L) or generalized forms(G) were identified.

Localized - < 30% of sites involved and CAL more than 1mm.

Generalized - > 30% of sites involved and CAL more than 1mm.

Subjects who were clinically healthy, with probing depth ≤ 3 mm and no clinical attachment loss were considered as healthy(H) subjects.

Probing depth (Grant 1965)⁴¹

Probing depth is measured from the gingival margin to the base of the pocket using a William's periodontal probe. The probe is passed under the gingiva along the circumference of the tooth. The probe inserted is always maintained parallel to the long axis of the tooth. Three measurements are made on the buccal aspect and three on the lingual aspect of each tooth – total of six sites per tooth.

Clinical attachment level (Carranza)⁸⁶

Clinical attachment level is measured from the cemento enamel junction to the base of the pocket using a William's periodontal probe.

When the gingival margin is located on the anatomic crown the level of attachment is determined by subtracting from the pocket depth, the distance from gingival margin to the cemento enamel junction. If both

are the same, the loss of attachment is zero. When the gingival margin coincides with the CEJ, the loss of attachment equals the pocket depth.

When the gingival margin is located apical to CEJ, the loss of attachment is greater than the pocket depth and therefore the distance between CEJ and gingival margin should be added to the pocket depth.

SPECIMEN COLLECTION

Patients were instructed to fast overnight before endoscopy.


Prior to endoscopic examination of the patients, plaque samples were collected. Dental plaque was removed from the tooth surfaces with a sterile periodontal curette. The sample was dispersed immediately into a vial containing 1ml of urea broth with phenyl red indicator to detect urease activity, and in another vial containing 0.2% sterile isotonic saline for culture.

Endoscopy was carried out with an Olympus fiber optic endoscope by a Gastroenterologist. Prior to specimen collection, the endoscope with biopsy forceps was rinsed thoroughly in water and soaked in 2% glutaraldehyde (CIDEX) for 20 minutes. The endoscope was thoroughly rinsed with sterile normal saline just before use. From each subject, four biopsy specimens were taken from antral mucosa 2 cm from the pylorus³⁵. The specimens were used for rapid urease test, culture, histopathology, Gram and Giemsa staining.

The specimens for rapid urease test were inoculated into vials containing urea broth immediately and the specimens for culture were inoculated into vials containing 0.2% sterile isotonic saline. The

specimens for histopathological examination were placed in 10% formalin. The specimens for culture were transported in an ice box to the laboratory and plated onto culture media within one hour of obtaining the specimen.

BIOPSY



BLOOD

With aseptic precautions, 2.5ml of venous blood was collected from each patient, the serum separated and stored at -20°C, till it was used for IgG antibody estimation.

PROCEDURE

RAPID UREASE TEST

To adapt to its special ecological niche, where the concentration of urea diffusing from blood to the gastric mucosa is low, *H.pylori* produces large amounts of urease. *H.pylori* urease also has the highest specific activity (36 ± 28 $\mu\text{mol/min mg}$ of protein among bacterial ureases *Mobley 1988*.⁸¹ The other urease positive bacteria present in the gastric mucosa i.e., streptococci and staphylococci produce a lower

amount of urease which does not interfere in a short – time detection (<2h), rendering the method specific to *H. pylori*.³⁵

In the oral cavity, urease positive microorganisms such as *Streptococcus vestibularis* and *Actinomyces viscosus* usually cannot give positive results within an hour. Hence rapid urease test has been used by several studies to detect the presence of *H.pylori* in dental plaque.

When a biopsy specimen or a plaque sample containing *H.pylori* is introduced into urea rich medium, the urease breaks the urea down into carbon dioxide and ammonia. The ammonium ion increases the pH, and a pH indicator eg.phenol red changes color in this case from yellow to red or violet.

UREA BROTH:

Stock solution A [1% phenol red solution (free acid) indicator]

1 gm of phenol red was dissolved in 32.5ml of 0.1 mol/l sodium hydroxide and made upto 100 ml with distilled water. The solution was autoclaved for 15 minutes at 121°C.

Stock solution B (10% urea solution)

10 gm of urea was dissolved in 100ml of sterile distilled water under sterile precautions. The pH of the solution was adjusted to 6.8 and dispensed in 0.5 ml aliquots in sterile vials.

Two drops of 1% phenol red (above) was added to each vial containing 0.5ml of 10% urea solution.

The specimens for the rapid urease test – antral biopsy specimen and the plaque sample were inoculated into two separate vials containing the urea broth. They were observed for color change from yellow to pink at room temperature every minute for the first 10 minutes, every 10 minutes for one hour and thereafter hourly for 8 hours and then at 24 hours. The change occurring within 2 hours denotes a positive reaction.³⁵

CULTURE⁶⁰

The antral biopsy specimens and the plaque samples, were transferred to the microbiology lab within 20 minutes at 2-8°C in an ice pack. The biopsy specimen was finely ground between two sterile frosted glass slides. Both the biopsy and the plaque samples were then inoculated onto selective – Campylobacter agar base – Skirrow's medium and non selective media – chocolate agar.

NON SELECTIVE MEDIA

Chocolate Agar

Peptone	:	1.0 gm
Meat extract	:	1.0gm
Sodium chloride	:	0.5gm
Agar agar	:	2 gm
Distilled water	:	100 ml
Defibrinated sheep blood	:	10%

SELECTIVE MEDIA

Skirrow's Campylobacter medium

- i) Campylobacter agar base : 39.5 gm
- ii) Distilled water : 1 liter
- iii) Skirrow's supplement : 1 vial consisting of
 - Vancomycin : 10 gm
 - Polymyxin B : 2500 IU
 - Trimethoprim : 5 ml
- iv Defibrinated sheep blood : 50 ml

The plates were incubated immediately at 37°C in a microaerophilic atmosphere in a candle jar or in a BOD (biological oxygen demand) incubator with Co₂ supply for 5 days.

Colony Characteristics

Chocolate agar plate in - Growth obtained from 2 antral
BOD incubator biopsies

The colonies showed the following morphology – circular, convex, translucent and glistening with slight hemolysis in blood agar.

The suspected colonies were confirmed by

- 1) **Gram stain** – gram negative curved bacilli were seen – spiral forms were less obvious.
- 2) **Oxidase test** – The suspected colony was streaked on the surface of an oxidase strip. An intense purple color within 5 seconds was noted as positive.

3) *Catalase test* – The suspected colony was introduced into hydrogen peroxide. Immediate production of gas bubbles was noted as positive.

4) *Urease test* – The suspected colony was introduced into 0.5ml of the urea broth. An instant color change from yellow to pink, which deepened in intensity, was noted as positive.

HISTOPATHOLOGY

The biopsy samples from the antrum were processed and stained with haematoxylin and eosin stain and examined for *Helicobacter pylori*.

CRUSH CYTOLOGY⁹⁷

The specimen in the transport medium was crushed between two sterile glass slides.

Gram stain

One slide was air dried and heat fixed. The slide was first covered with methyl violet for one minute. The excess stain was poured off, gram's iodine added and after 1 minute washed and decolorized with acetone for 2-3 seconds. The acetone was washed off with water and counter stained with dilute carbol fuschin for one minute. The slide was then washed with water, blotted dry and observed under oil immersion objective. The smear was examined for the presence of gram negative helical bacteria.

Giemsa stain

The other slide was air dried and fixed with methanol for 3 minutes. 2 – 3 drops of undiluted Giemsa stain (Qualigenas, Glaxo) was added and kept for 5 minutes. The smear was then washed with water, blotted dry and seen under oil immersion objective. The organism appeared deep purple with the typical gull-wing morphology.

SEROLOGY

The serological detection of IgG antibodies to cellular components of H.pylori was done, using the **EUROIMMUNO ELISA Kit**.

Antigen: The antigen source is provided by the strain Lior 1, (Brussels) “of Helicobacter pylori. The cultured bacteria have been disrupted in alkaline buffer.” The used antigen mixture contains all significant proteins as verified by SDS (sodium dodecyl sulphate) polyacrylamide gel electrophoresis.

Contents of the test kit:

1. Micro plate wells coated with antigens
2. Calibrator 1
200 RU/ml (IgG, Human), ready for use
3. Calibrator 2
20 RU/ml (IgG, Human), ready for use
4. Calibrator 3
2 RU/ml (IgG, Human), ready for use
5. Positive control
(IgG, Human), ready for use
6. Negative Control
(IgG, Human), ready for use
7. Enzyme conjugate
Peroxidase – Labeled anti – human IgG (rabbit), ready for use.
8. Sample buffer ready for use
9. Wash buffer
10 xs concentrate
10. Chromogen / substrate solution

TMB / H₂O₂ ready for use

11. Stop solution

0.5 sulphuric acid, ready for use

12. Test instruction

13. Protocol with target values

Serology was done for 110 patients. 3 calibrated standards provided by the manufacturer and serum samples of all the cases and controls were included.

Method:

- Serum samples were diluted 1: 101 before assay (10 µl of serum was diluted with 1ml of sample diluents).
- 100 µl of each calibrated standard or diluted sample was dispensed into the wells.
- The plate was incubated for 30 minutes at room temperature.
- The wells were washed thoroughly thrice using wash buffer, the micro plate was blotted on absorbent paper.
- 100 µl conjugate consisting of rabbit antihuman IgG conjugated with Horse radish peroxidase was added immediately into each well.
- The plate was incubated for 30 minutes at room temperature.
- Following incubation, the plate was washed 3 times with wash buffer.
- 100µl of chromogen / substrate solution was added into each well. The plate was incubated for 15 minutes. Green color developed in the wells.
- 100µl of stop solution (0.05m sulphuric acid) was added to each well in the same as the chromogen substrate to allow equal reaction times.
- Green color change to yellow on addition of the stop solution.

- The optical density was read at 450 nm in a micro plate reader within 30 minutes of adding the stop solution.

CALCULATION OF RESULTS

The optical density of each calibrator was plotted against its concentration and a curve was drawn through the points. The unknowns were read off the curve.

The kit recommends interpreting results as follows:

< 16 RU/ml –Negative

≥ 16 to <22 RU/ml – Borderline

≥ 22 RU/ml – Positive

ARMAMENTARIUM

Periodontal examination

Mouth mirror

William's periodontal probe

Explorer

Tweezers

Mask

Head cap

Gloves

Periodontal curettes

Sterile aliquots containing saline

Rapid Urease Test

Test tube

Beaker

Conical flask

Dropper

Forceps

Sterile aliquots

Culture

Culture (media plates)

Bacteriological loop

Sterile stick

Petri disk

Test tubes

Sterile aliquots

Frosted glass slides

Bunsen burner

Carbon dioxide incubator

Candle jar

Ice base

Histopathology

Plane glass slide

Microscope

ELISA

Test tubes

Wooden Rack

Measuring Jar

Bowls

Autoclaved plastic tips

ELISA washer

ELISA reader



Photograph No.1: Good oral hygiene in H.pylori positive patient



Photograph No.2: Fair oral hygiene in H.pylori positive patient



Photograph No.3: Poor oral hygiene in H.pylori positive patient



Photograph No.4: Periodontitis in H.pylori positive patient



Photograph No.5: Armamentarium for dental plaque and blood sampling



Photograph No.6: Pocket depth > 5mm in H.pylori positive patient



Photograph No.7: Aliquots with urea broth and saline for RUT and sample collection



Photograph No.8: Transport for samples



Photograph No.9: Collection of dental plaque sample



Photograph No. 10: Endoscope and biopsy forceps for antral biopsy



Photograph No. 11: Antral biopsy obtained during endoscopy by Gastroenterologist



Photograph No.12: Antral biopsy taken from the biopsy forceps



Photograph No.13: Collection of blood sample for ELISA



Photograph No.14: Armamentarium for culture (left), streaking of sample onto culture plate (right)




Photograph No. 15: Electron Microscope



Photograph No. 16: ELISA kit for Anti-H.pylori (IgG)



Photograph No. 17: Serum samples for ELISA



Photograph No. 18: Microplate wells coated with antigen

STATISTICAL ANALYSIS

The statistical analysis was done using the computer software program SPSS version 12.

Mean and Standard Deviation were estimated for different variables in each study group.

Normality of the data was tested in each group by using Kolmogorov Smirnov test.

Mean values were compared between two study groups by using either *Student's Independent t-test* or *Mann-Whitney U-Test*.

Pearson's chi-square test was used to compare the proportions two study groups.

In the present study, $p < 0.05$ was considered as the level of significance.

Statistical Formulae Used For Data Analysis

Pearson's Chi-square Test

The formula used was

$$\chi^2 = \sum_{i=1}^n \left\{ \frac{(O - E)^2}{E} \right\}$$

Where O = Observed frequency in a cell

E = Expected frequency in a cell

Pearson's Chi-square Test with Yates' Continuity Correction

The formula used was

$$\chi^2 = \sum_{i=1}^n \frac{[O - E - \frac{1}{2}]^2}{E}$$

Where O = Observed frequency in a cell

E = Expected frequency in a cell

Mann-Whitney U-Test

The formula is

$$Z = \frac{\left[T - \frac{n_1(n_1 + n_2 + 1)}{2} \right]}{\sqrt{\frac{n_1 n_2 (n_1 + n_2 + 1)}{12}}}$$

Where n1 and n2 are the sample sizes in Group I and Group II respectively.

T is the sum of the ranks for the n1 observations.

Student's Independent t-test

The formula used is

$$t = (\bar{X}_1 - \bar{X}_2) / \sqrt{S_p^2 / n_1 + S_p^2 / n_2}$$

where $S_p^2 = [(n_1 - 1)S_1^2 + (n_2 - 1)S_2^2] / (n_1 + n_2 - 2)$

\bar{X}_1 and \bar{X}_2 are the sample means;

RESULTS

One hundred and ten subjects were included in the study. The subjects were categorized into two groups – cases (n = 55) and controls (n = 55).

Table I & II shows the master chart of cases and controls with the variables, clinical parameters and tests done in both groups in the study.

Table III & Fig I shows the comparison of gender between cases and controls. Distribution of gender in both groups was similar.

Table IV & Fig II shows the comparison of handling of animals between cases and controls. 20 of 55 cases (36.4%) had contact with animals compared to only 4 of 55 controls (7.3%). This difference was found to be significant with a P value of 0.001.

Table V & Fig III shows the comparison of socioeconomic status between cases and controls. There were more patients of lower socioeconomic status in both cases (n=42/55), and controls (n=41/55) but it was not found to be statistically significant (P= 1.00).

Table VI shows the comparison of mean values between cases and controls. The mean age in controls (48 ± 12.5) was higher than cases (44.4 ± 15.2). However there was no significant difference in mean age between cases and controls. The mean socio-economic status score in controls (10.0 ± 4.2) was higher than in cases (9.4 ± 4.1). However there was no significant difference in mean SES score between cases and controls (P = 0.27). The mean ELISA (RU/ml) score in cases (159 ± 32) was significantly higher than in controls (54 ± 36) P (< 0.0001). The mean OHI score in cases (2.05 ± 1.16) was higher than in controls

(1.98 ± 1.07). However, there was no significant difference in OHI scores between cases and controls ($P=0.75$).

Table VII shows the comparison of smoking habit and alcohol consumption between cases and controls. The proportion of current smokers in cases (38.2%) was slightly higher than controls (29.1). However there was no significant difference in the proportion of current smokers between the two study groups ($P = 0.59$). The proportion of people currently having the habit of alcohol consumption in cases (41.8%) was slightly lower than controls (43.6%). However there was no significant difference between cases and controls ($P = 0.67$).

Table VIII & Fig IV shows the comparison of OHI score between cases and controls. Among cases 19 subjects (34.5%) had good oral hygiene, 25(45.5%) had fair and 11 (20.0%) had poor oral hygiene. Among controls 17(30.9%), 28 (50.9%), and 10 (18.2%) had good, fair and poor oral hygiene respectively. The observed difference in the oral hygiene status between the two groups was not found to be statistically significant ($P = 0.85$).

Table IX & Fig V shows the comparison of periodontal disease and pocket depth $> 5\text{mm}$ between cases and controls. 27 (49.1%) subjects among cases had periodontal disease with pocket depth of $\geq 5\text{mm}$ in at least one site compared to only 8 (14.5%) subjects among controls. This difference was found to be statistically significant ($P < 0.001$).

Table X & Fig VI shows the comparison of RUT (P) and culture (P) between cases and controls. 48 cases (87.3%) had a positive rapid urease test compared to 29 (52.7%) subjects among controls. This difference was found to be statistically significant with P value of $<$

0.001. H.pylori could not be cultured from dental plaque in any of the 55 cases or controls.

Table XI shows the comparison of RUT (B), culture (B), Gram and Giemsa stain between cases and controls. On antral biopsy specimens, all 55 cases (100%) had positive rapid urease test, while 55 controls had negative RUT. 2 cases tested positive on culture, 40 cases (72.7%) positive on Gram staining and 50 cases (90.9%) positive on Giemsa staining for H.pylori.

Results of histopathological examination of antral biopsy samples revealed 40 cases with chronic active gastritis, 4 cases with features suggestive of malignancy (adenocarcinoma), 2– cases with acute gastritis, 5 cases of atrophic gastritis and 4 cases of intestinal metaplasia. Among controls the picture was that of chronic nonactive gastritis or lymphocytic infiltration

Table XII shows the results of multiple logistic regression analysis. The variables found to be statistically significant in the university analysis-contact with animals; periodontal disease status with pocket depth of ≥ 5 mm and RUT (P) positive were included in the multivariate logistic regression analysis. In the multivariate analysis, it was found that the observed difference in the above three variables between the two groups were statistically significant.

Tables XIII and IV shows the association of OHI and Periodontal disease with RUT (P) between cases and controls. The results showed that the association of OHI and Periodontal disease with RUT(P) in cases was not significant, but among controls the association of periodontal disease with RUT(P) was significant($P=0.005$).

TABLE –I MASTER CHART-CASES

Code No	Age	Sex	Smoker			Alcohol			Contact with Animals	SES Score	SES Status	OHI Score	OHI Status	Periodontal disease status	Pocket depth > 5mm / site	RUT		Culture		ELISA (RU/ml)	Gram Stain	Giemsa stain
			Current	Ex	No	Current	Past	No								P	B	P	B			
001	35	M	✓			✓			No	7	(U) Lower	2.0	Fair	H	-	+	+	-	-	190	+	+
002	67	M	✓			✓			Yes Cats	6	(U) Lower	2.9	Fair	G	+	+	+	-	-	136	+	+
003	75	M		✓			✓		No	16	(U) Middle	3.2	Poor	G	+	+	+	-	-	140	+	+
004	45	M	✓			✓			No	11	(L) Middle	3.2	Poor	G	+	-	+	-	-	168	+	+
005	21	M			✓			✓	Yes Dogs Goats	8	(U) Lower	1.3	Good	H	-	+	+	-	-	> 200	+	+
006	67	M		✓			✓		Yes Cows Dogs	12	(L) Middle	1.0	Good	H	-	+	+	-	-	114	-	+
007	66	M			✓	✓		✓	No	11	(L) Middle	2.2	Fair	H	-	+	+	-	-	136	+	+
008	39	M		✓		✓			No	7	(U) Lower	1.3	Good	H	-	+	+	-	-	138	+	+
009	35	M		✓		✓			No	6	(U) Lower	2.1	Fair	H	-	+	+	-	-	135	+	+
010	45	M	✓			✓			No	7	(U) Lower	2.9	Fair	H	-	+	+	-	-	> 200	+	+
011	30	M	✓			✓			Yes Cats	16	(U) Middle	3.6	Poor	G	+	+	+	-	-	> 200	+	+
012	51	M		✓		✓			No	10	(U) Lower	2.0	Fair	L	+	+	+	-	-	114	+	+
013	52	M	✓			✓			No	8	(U) Lower	3.3	Poor	G	+	+	+	-	-	188	+	+
014	33	F			✓			✓	Yes Cats	9	(U) Lower	1.2	Good	H	-	+	+	-	-	> 200	+	+
015	70	M		✓			✓		Yes Goats	19	(U) Middle	2.1	Fair	L	+	+	+	-	-	> 200	+	+
016	30	M			✓			✓	No	16	(U) Middle	0.6	Good	H	-	+	+	-	-	> 200	+	+
017	45	F			✓			✓	No	9	(U) Lower	2.7	Fair	H	-	+	+	-	-	164	-	+
018	33	F			✓			✓	No	5	(U) Lower	0.8	Good	H	-	-	+	-	-	110	+	+
019	44	M		✓			✓		No	8	(U) Lower	2.6	Fair	G	+	+	+	-	-	198	-	+
020	40	M	✓					✓	Yes Goats	6	(U) Lower	3.4	Poor	G	+	+	+	-	-	90	+	+
021	69	M		✓			✓		No	14	(U) Middle	2.0	Fair	G	+	+	+	-	-	190	+	+
022	70	M			✓	✓			Yes Cats	5	Lower	2.0	Fair	G	+	-	+	-	-	170	-	+
023	47	M		✓		✓			Yes Goats	8	(U) Lower	2.2	Fair	L	+	-	+	-	-	96	+	+
024	18	F			✓			✓	Yes Dogs	4	Lower	0.46	Good	H	-	-	+	-	-	170	+	+
025	21	F							Yes Goats	10	(U) Lower	0.46	Good	H	-	-	-	-	-	> 200	+	+

Code No	Age	Sex	Smoker			Alcohol			Contact with Animals	SES Score	SES Status	OHI Score	OHI Status	Periodontal disease status	Pocket depth > 5mm / site	RUT		Culture		ELISA (RU/ml)	Gram Stain	Giemsa stain
			Current	Ex	No	Current	Past	No								P	B	P	B			
026	50	M	✓			✓			Yes Hens	9	(U) Lower	4.6	Poor	G	+	+	+	-	-	138	+	+
027	62	M	✓			✓			Yes Cows	10	(U) Lower	2.0	Fair	G	+	+	+	-	-	156	+	+
028	44	M	✓					✓	No	7	(U) Lower	3.0	Fair	H	-	+	+	-	-	130	+	+
029	33	F			✓			✓	Yes Dogs	3	Lower	0.6	Good	H	-	+	+	-	-	180	+	+
030	65	M	✓			✓			No	16	(U) Middle	4.0	Poor	G	+	+	+	-	-	136	+	+
031	57	M	✓			✓			Yes Cows	8	(U) Lower	3.0	Fair	L	+	+	+	-	-	164	+	+
032	46	M			✓	✓			Yes Goats	9	(U) Lower	1.8	Fair	L	+	+	+	-	-	198	+	+
033	47	M	✓			✓			Yes Cats	14	(L) Middle	2.7	Fair	G	+	+	+	-	-	150	+	+
034	65	M		✓			✓		No	24	(U) Middle	2.4	Fair	L	+	+	+	-	-	167	+	+
035	36	F			✓			✓	No	8	(U) Lower	0.7	Good	H	-	+	+	-	-	158	+	+
036	35	F			✓			✓	No	7	(U) Lower	1.7	Fair	H	-	+	+	-	-	118	-	+
037	19	M			✓			✓	No	5	(U) Lower	1.1	Good	H	-	+	+	-	-	174	+	+
038	55	M		✓				✓	No	10	(U) Lower	5.0	Poor	G	+	+	+	-	-	136	+	+
039	47	M	✓					✓	Yes Cows	8	(U) Lower	2.1	Fair	H	-	+	+	-	-	170	+	+
040	35	M			✓			✓	No	7	(U) Lower	0.6	Good	H	-	+	+	-	-	168	+	+
041	50	M	✓					✓	No	11	(L) Middle	1.8	Fair	G	+	+	+	-	-	170	+	+
042	38	M			✓	✓			No	9	(U) Lower	0.9	Good	H	-	+	+	-	-	168	+	+
043	35	F	✓					✓	No	8	(U) Lower	3.1	Poor	G	+	+	+	-	-	106	-	+
044	35	M	✓			✓			No	9	(U) Lower	3.2	Poor	L	+	+	+	-	-	150	+	+
045	30	M	✓			✓			No	10	(U) Lower	1.6	Fair	H	-	+	+	-	-	135	-	+
046	35	F			✓			✓	No	8	(U) Lower	1.2	Good	H	-	+	+	-	-	140	-	-
047	45	M	✓			✓			Yes Dogs Goats	10	(U) Lower	2.3	Fair	L	+	+	+	-	-	155	-	-
048	46	F			✓			✓	No	8	(U) Lower	1.2	Fair	L	+	+	+	-	-	160	-	+
049	67	M	✓			✓			Yes Dogs	19	(U) Middle	2.0	Fair	G	+	+	+	-	-	130	-	-
050	26	F			✓			✓	No	9	(U) Lower	1.0	Good	H	-	+	+	-	-	146	-	-
051	17	F			✓			✓	No	6	(U) Lower	0.6	Good	H	-	+	+	-	-	127	-	+
052	44	F			✓			✓	No	6	(U) Lower	0.8	Good	H	-	+	+	-	-	137	-	+
053	65	M		✓			✓		No	9	(U) Lower	5.0	Poor	G	+	+	+	-	-	200	+	-
054	35	F			✓			✓	No	4	(U) Lower	0.6	Good	H	-	+	+	-	-	125	-	+
055	33	M	✓			✓			No	9	(U) Lower	1.0	Good	H	-	-	+	-	--	136	-	+

H-Healthy; G- Generalized chronic periodontitis; L- Localized chronic periodontitis

Table –II MASTER CHART- CONTROLS

Code No	Age	Sex	Smoker			Alcohol			Contact with Animals	SES Score	SES Status	OHI Score	OHI Status	Periodontal disease status	Pocket depth > 5mm / site	RUT		Culture		ELISA (RU/ml)	Gram Stain	Giemsa stain
			Current	Ex	No	Current	Past	No								P	B	P	B			
001	46	M	✓					✓	No	12	(L) Middle	2.9	Fair	H	-	-	-	-	-	12	-	-
002	46	M	✓					✓	No	6	(U) Lower	4.3	Poor	H	-	-	-	-	-	14	-	-
003	65	M	✓			✓			No	8	(U) Lower	3.1	Poor	H	-	-	-	-	-	22	-	-
004	45	M		✓				✓	No	8	(U) Lower	1.1	Good	H	-	+	-	-	-	19	-	-
005	37	M	✓				✓		No	10	(U) Lower	0.3	Good	H	-	+	-	-	-	26	-	-
006	65	M			✓	✓			No	10	(U) Lower	2.5	Fair	G	+	+	-	-	-	32	-	-
007	47	F			✓			✓	No	4	(U) Lower	0.6	Good	H	-	+	-	-	-	12	-	-
008	40	F			✓			✓	No	9	(U) Lower	0.6	Good	H	-	-	-	-	-	70	-	-
009	43	M	✓			✓			No	9	(U) Lower	4.0	Poor	G	+	+	-	-	-	48	-	-
010	26	M			✓			✓	No	16	(U) Middle	0.9	Good	H	-	-	-	-	-	48	-	-
011	30	M			✓	✓			Yes Cows	10	(U) Lower	2.0	Fair	H	-	+	-	-	-		-	-
012	40	F							No	6	(U) Lower	3.6	Poor	G	+	+	-	-	-	78	-	-
013	57	M	✓			✓			No	18	(U) Middle	1.8	Fair	H	-	-	-	-	-	62	-	-
014	50	M	✓			✓			No	9	(U) Lower	2.0	Fair	H	-	+	-	-	-	70	-	-
015	58	M		✓				✓	Yes Cows	10	(U) Lower	2.4	Fair	H	-	+	-	-	-		-	-
016	70	M							No	8	(U) Lower	1.8	Fair	H	-	-	-	-	-	78	-	-
017	50	M	✓					✓	Yes Dogs, Cows, Hens	8	(U) Lower	2.4	Fair	G	+	+	-	-	-	78	-	-
018	40	F			✓			✓	No	14	(U) Middle	0.6	Good	H	-	-	-	-	-	60	-	-
019	36	M			✓	✓			No	8	(U) Lower	0.4	Good	H	-	+	-	-	-	94	-	-
020	47	M			✓	✓			No	16	(U) Middle	3.2	Poor	H	-	-	-	-	-	24	-	-
021	50	F			✓			✓	No	5	(U) Lower	0.9	Good	H	-	+	-	-	-	20	-	-
022	70	F			✓			✓	No	2	Lower	3.4	Poor	G	+	+	-	-	-	14	-	-
023	46	M			✓	✓			No	9	(U) Lower	2.6	Fair	H	-	+	-	-	-	92	-	-
024	46	M	✓			✓			No	18	(U) Middle	2.0	Fair	H	-	-	-	-	-	100	-	-
025	76	M		✓			✓		No	15	(L) Middle	2.8	Fair	H	-	-	-	-	-	18	-	-
026	55	M	✓	□		□	✓		No	8	(U) Lower	0.6	Good	H	-	+	-	-	-	74	-	-
027	35	M	✓			✓			No	15	(L) Middle	3.2	Poor	L	+	+	-	-	-	50	-	-
028	38	M	✓			✓			No	10	(L) Lower	2.2	Fair	H	-	-	-	-	-	104	-	-
029	70	M			✓	✓			No	9	(L) Lower	3.0	Fair	H	-	+	-	-	-	60	-	-
030	39	M		✓		✓			No	9	(L) Lower	1.6	Fair	H	-	-	-	-	-	18	-	-

H-Healthy; G- Generalized chronic periodontitis; L- Localized chronic periodontitis

Code No	Age	Sex	Smoker			Alcohol			Contact with Animals	SES Score	SES Status	OHI Score	OHI Status	Periodontal disease status	Pocket depth > 5mm / site	RUT		Culture		ELISA (RU/ml)	Gram Stain	Giemsa stain
			Current	Ex	No	Current	Past	No								P	B	P	B			
031	65	M		✓			✓		No	9	(U) Lower	2.3	Fair	H	-	+	-	-	-	78	-	-
032	46	M	✓			✓			No	18	(U) Middle	1.5	Fair	H	-	-	-	-	-	120	-	-
033	65	M	✓			✓			No	13	(U) Middle	2.2	Fair	H	-	-	-	-	-	72	-	-
034	30	M			✓	✓			No	9	(U) Lower	1.8	Fair	H	-	+	-	-	-	18	-	-
035	48	M	✓			✓			No	6	(U) Lower	1.2	Good	H	-	+	-	-	-	98	-	-
036	45	M			✓			✓	No	10	(U) Lower	1.5	Fair	H	-	-	-	-	-	3	-	-
037	40	M			✓	✓			No	8	(U) Lower	2.0	Fair	H	-	-	-	-	-	6	-	-
038	38	M		✓				✓	No	10	(U) Lower	4.0	Poor	H	-	+	-	-	-	8	-	-
039	25	M			✓			✓	No	25	(U) Middle	0.3	Good	H	-	-	-	-	-	18	-	-
040	59	M	✓			✓			No	15	(L) Middle	1.4	Fair	H	-	-	-	-	-	22	-	-
041	41	M							No	11	(L) Middle	0.6	Good	H	-	-	-	-	-	34	-	-
042	62	M		✓				✓	No	15	(L) Middle	2.2	Fair	H	-	+	-	-	-	64	-	-
043	68	M		✓		✓			No	8	(U) Lower	1.9	Fair	H	-	-	-	-	-	20	-	-
044	60	M		✓		✓			No	10	(U) Lower	2.8	Fair	H	-	+	-	-	-	58	-	-
045	42	M	✓			✓			Yes Cows	10	(U) Lower	2.0	Fair	L	+	+	-	-	-	96	-	-
046	43	F			✓			✓	No	5	(U) Lower	1.2	Good	H	-	-	-	-	-	23	-	-
047	43	F	✓			✓			No	10	(U) Lower	2.2	Fair	H	-	+	-	-	-	40	-	-
048	27	F			✓			✓	No	9	(U) Lower	0.6	Good	H	-	+	-	-	-	16	-	-
049	39	F			✓			✓	No	9	(U) Lower	3.1	Poor	H	-	-	-	-	-	26	-	-
050	41	F			✓			✓	No	9	(U) Lower	0.9	Good	H	-	+	-	-	-	18	-	-
051	45	F			✓			✓	No	4	(U) Lower	2.0	Fair	H	-	-	-	-	-	106	-	-
052	42	F			✓			✓	No	5	(U) Lower	0.4	Good	H	-	+	-	-	-	112	-	-
053	45	F			✓			✓	No	10	(U) Lower	0.9	Good	H	-	-	-	-	-	14	-	-
054	60	F			✓			✓	No	8	(U) Lower	2.6	Fair	H	-	-	-	-	-	126	-	-
055	67	F			✓			✓	No	4	(U) Lower	4.1	Poor	G	+	+	-	-	-	53	-	-

TABLE-III

COMPARISON OF GENDER BETWEEN CASES AND CONTROLS

Variable	Category	Cases [n=55]		Controls [n=55]		P – value*
		No.	%	No.	%	
Gender	Male	16	29.1	16	29.1	1.00(NS)
	Female	39	70.9	39	70.9	

FIGURE I

DISTRIBUTION OF GENDER IN CASES AND CONTROLS




TABLE-IV
COMPARISON OF HANDLING OF ANIMALS BETWEEN CASES AND CONTROLS

Variable	Category	Cases [n=55]		Controls [n=55]		P – value*
		No.	%	No.	%	
Contact with animals	No	35	63.6	51	92.7	0.001 (Sig.)
	Yes	20	36.4	4	7.3	
Type of animal	Cat	6	30.0	0	0.0	0.07 (N.S)
	Dog	3	15.0	0	0.0	
	Cow	3	15.0	3	75.0	
	Goat	3	15.0	0	0.0	
	Hen	1	5.0	0	0.0	
	Dog, Cow, Goat	0	0.0	1	25.0	
	Dog, Goat	3	15.0	0	0.0	
	Cow, Dog	1	5.0	0	0.0	

FIGURE II
COMPARISON OF HANDLING OF ANIMALS BETWEEN CASES AND CONTROLS




TABLE-V
COMPARISON OF SOCIOECONOMIC STATUS BETWEEN CASES AND CONTROLS

Variable	Category	Cases [n=55]		Controls [n=55]		P – value*
		No.	%	No.	%	
SES	Lower	42	76.4	41	74.5	1.00 (N.S.)
	Middle	13	23.6	14	25.5	

FIGURE III
COMPARISON OF SOCIOECONOMIC STATUS BETWEEN CASES AND CONTROLS




TABLE VI
COMPARISON OF MEAN VALUES BETWEEN CASES AND CONTROLS

Variable	Group	Mean \pm S.D.	P-value*
Age	Cases	44.4 \pm 15.2	0.17 (N.S.)
	Controls	48.2 \pm 12.5	
OHI Score	Cases	2.05 \pm 1.16	0.75 (N.S.)
	Controls	1.98 \pm 1.07	
ELISA	Cases	159 \pm 32	<0.0001 (Sig.)
	Controls	54 \pm 36	
SES Score	Cases	9.4 \pm 4.1	0.27** (N.S.)
	Controls	10.0 \pm 4.2	

*Student's independent t-test was used to calculate the P-value.

** Mann-Whitney U-test was used to calculate the P-value.

TABLE-VII
**COMPARISON OF SMOKING HABIT AND ALCOHOL CONSUMPTION
BETWEEN CASES AND CONTROLS**

Variable	Category	Cases [n=55]		Controls [n=55]		P – value*
		No.	%	No.	%	
Smoker	Non-smoker	22	40.0	26	47.3	0.59(NS)
	Current	21	38.2	16	29.1	
	Ex- smoker	12	21.8	13	23.6	
Alcohol	Not at all	24	43.6	26	47.3	0.67(NS)
	Current	23	41.8	24	43.6	
	Past	8	14.5	5	9.1	

TABLE-VIII

COMPARISON OF OHI SCORE BETWEEN CASES AND CONTROLS

Variable	Category	Cases [n=55]		Controls [n=55]		P – value*
		No.	%	No.	%	
OHI- SCORE	Poor	11	20.0	10	18.2	0.85 (N.S.)
	Fair	25	45.5	28	50.9	
	Good	19	34.5	17	30.9	

FIGUREIV

**COMPARISON OF OHI SCORE BETWEEN CASES AND
CONTROLS**




TABLE-IX

**COMPARISON OF PERIODONTAL DISEASE AND POCKET DEPTH
≥ 5MM BETWEEN CASES AND CONTROLS**

Variable	Category	Cases [n=55]		Controls [n=55]		P – value*
		No.	%	No.	%	
Periodontal disease	Healthy	28	50.9	47	85.5	<0.0001 (Sig.)
	Periodontitis	27	49.1	8	14.5	
Pocket depth > 5mm / site	No	28	50.9	47	85.5	<0.0001 (Sig.)
	Yes	27	49.1	8	14.5	

FIGURE V

**COMPARISON OF PERIODONTAL DISEASE BETWEEN CASES AND
CONTROLS**




TABLE-X

COMPARISON OF RUT (P) AND CULTURE (P) BETWEEN CASES AND CONTROLS

Variable	Category	Cases [n=55]		Controls [n=55]		P – value*
		No.	%	No.	%	
RUT P	No	7	12.7	26	47.3	<0.0001 (Sig.)
	Yes	48	87.3	29	52.7	
Culture P	Negative	55	100.0	55	100.0	-
	Positive	0	0.0	0	0.0	

FIGURE VI

COMPARISON OF RUT (P) BETWEEN CASES AND CONTROLS




TABLE-XI
COMPARISON OF RUT (B), CULTURE (B), GRAM STAIN AND
GIEMSA STAIN BETWEEN CASES AND CONTROLS

Variable	Category	Cases [n=55]		Controls [n=55]		P – value*
		No.	%	No.	%	
RUT B	No	0	0.0	55	100.0	<0.0001 (Sig.)
	Yes	55	100.0	0	0.0	
	Positive	0	0.0	0	0.0	
Culture B	Negative	55	100.0	55	100.0	<0.0001 (Sig.)
	Positive	2	0.0	0	1.8	
Gram Stain	Negative	15	27.3	55	100.0	<0.0001 (Sig.)
	Positive	40	72.7	0	0.0	
Giemsa stain	Negative	5	9.1	55	100.0	<0.0001 (Sig.)
	Positive	50	90.9	0	0.0	

TABLE XII
RESULTS OF MULTIPLE LOGISTIC REGRESSION ANALYSIS

Independent Variable	Odds Ratio [95% CI]	P-value*
Contact with animals	6.13 [1.70 to 22.14]	0.006 (Sig.)
Periodontal disease Status	2.89 [1.06 to 7.89]	0.04 (Sig.)
RUT P	5.05 [1.71 to 14.89]	0.003 (Sig.)

TABLE-XIII

**ASSOCIATION OF OHI AND PERIODONTAL DISEASE WITH RUT P
IN CASES**

Variable	Category	RUT P - No [n=7]		RUT P - Yes [n=48]		P – value*
		No.	%	No.	%	
OHI-SCORE	Poor	1	14.3	10	20.8	0.40 (N.S.)
	Fair	2	28.6	23	47.9	
	Good	4	57.1	15	31.3	
Periodontal disease	Healthy	4	57.1	24	50.0	1.00** (N.S.)
	Periodontitis	0	0.0	27	53.8	

* Pearson's Chi-square test was used to calculate the P-value.

** Fisher's Exact Test (2 –tailed) was used to calculate the P-value.


TABLE-XIV

**ASSOCIATION OF OHI AND PERIODONTAL DISEASE WITH RUT P
IN CONTROLS**


Variable	Category	RUT P - No [n=26]		RUT P - Yes [n=29]		P – value*
		No.	%	No.	%	
OHI-SCORE	Poor	4	15.4	6	20.7	0.63 (N.S.)
	Fair	15	57.7	13	44.8	
	Good	7	26.9	10	34.5	
Periodontal disease	Healthy	26	100.0	21	72.4	0.005** (Sig.)
	Periodontitis	0	0.0	8	27.6	

* Pearson's Chi-square test was used to calculate the P-value.


** Fisher's Exact Test (2 –tailed) was used to calculate the P-value.




Photograph No.19: RUT positive in dental plaque and antral biopsy samples




Photograph No. 20: Culture positive for H.pylori in antral biopsy



Photograph No. 21: Giemsa stain showing H.pylori in antral biopsy specimen



Photograph No. 22: Histopathological features of chronic gastritis showing chronic lymphocytic infiltration



Photograph No.23: Microplate wells after ELISA reaction



Photograph No. 24: Microplate wells in ELISA reader

DISCUSSION

Periodontal disease may be related to a number of systemic diseases including increased incidence of atherosclerosis, coronary heart disease, stroke, diabetes mellitus, pre-term low birth weight delivery and respiratory diseases (*Mealey BL 1999*)⁷⁶.

*Robinson and Moynihan in 1904*⁷⁴ suggested that oral sepsis might play a part in the pathogenesis of gastric ulcers. The discovery of *Helicobacter pylori* and the acceptance of its role in gastric pathophysiology represented a fundamental change in the understanding of gastro duodenal disease (*Martin J.M. Buckley and Colm A.O' Morain 1998*).⁷³ It is now acknowledged worldwide that type B gastritis, duodenal ulceration, gastric ulceration, gastric adenocarcinoma and gastric mucosa associated lymphoid tissue lymphoma are in part infectious diseases.⁷³ Antibiotic based regimens are now the recommended treatment for peptic ulceration.⁷³

Recently attention has been focused on the importance of dental plaque in harbouring *H.pylori* and its role in the epidemiology of *H.pylori* infection.

Dental plaque has been defined as the diverse community of micro-organisms found on the tooth surface as a biofilm embedded in an extracellular matrix of polymers of host and microbial origin (*Marsh P.D. 2004*).⁶⁸ Dental plaque typically adheres to supragingival and subgingival tooth surfaces and it will quickly form in the absence of good oral hygiene measures.

Chronic periodontitis is the most prevalent form of periodontitis and occurs in response to chronic plaque and calculus accumulation (*Carranza*).⁸⁷

This study was undertaken i) to determine whether dental plaque harbors *H.pylori* and may act as a reservoir for the organism and ii) to determine the relation between oral hygiene status, periodontal disease status and *H.pylori* gastritis.

Patients with active bleeding ulcers were excluded from the study, because the rapid urease test (RUT) lacks sensitivity in *Helicobacter pylori* diagnosis when peptic ulcer disease presents with bleeding (*Lee J. M. et al 2000*).⁵⁶ Thus patients on chronic use of NSAIDS were also excluded, as NSAIDS are related to increased incidence of bleeding ulcers (*Francis Megraud and Philippe Lehours 2007*).³⁵

The major period of *H.pylori* acquisition is in childhood (*Roger. A. Feldman et al 1998*).⁹⁹ Once *H.pylori* is acquired, it persists into old age, unless eradicated by treatment (*Dowsett. S.A. M.J. Kowolik 2003*).²⁷ Hence the age group in the present study had a wide range of 17-76 years.

Seroprevalence of *H.pylori* infection is often similar in males and females(*Gasbarrini G. et al 1995*).³⁸ The gender distribution of cases and controls in the present study was similar indicating that there was little influence of these variables on the disease status of the two groups (Table III).

Lower socioeconomic status and or a low level of education are associated with an increase in the prevalence of *H.pylori* infection

(*Malathy HM et al 1996*).⁶³ As the socioeconomic status of individuals and countries has risen, the prevalence in younger generations has declined (*Haruma et al 1997*).⁴⁵ However in developing countries, socioeconomic status and sanitary conditions have improved even more slowly, which is thought to account for the continuing high rates of infection in young people. In the present study although there were more subjects of lower socioeconomic status, there was no significant difference in the mean SES score between cases and controls (Tables V and VI).

Handling of animals by the subjects in the two groups was also evaluated (Table IV). The proportion of subjects who had contact with animals among cases (36.4%) is significantly higher than among controls (7.3%), with a P value of 0.001 and an odds ratio of 6.13 (by multiple logistic regression analysis) 95% CI: 1.70 to 22.14 (Table XII). Animals have been shown to harbor *H.pylori*. Cats are found with *H.pylori* in nature and could be a reservoir for human infection (*Fox JG. 1995*).³⁴ In the present study, the subjects had contact with animals like cats, dogs, hens, goats and cows. However the type of animal the subjects had contact with was not statistically significant, $P = 0.07$ (Table IV). This may be due to the small sample size and studies on larger population may help identify the animal with high risk for acquiring *H.pylori*.

The high degree of association between handling of animals and *H.pylori* infection in the present study, suggests that animals may play an important role in the transmission of *H.pylori* infection and that handling of animals may be an important risk factor for developing *H.pylori* infection.

The oral hygiene status of patients was examined using the Simplified Oral Hygiene Index of Greene and Vermilion 1964 in the present study. The mean OHI score in cases is higher than controls. However there is no significant difference in OHI score between cases and controls ($P = 0.75$) (Table V). This is similar to the study by **Pradeep et al 2006**,⁹⁴ wherein the OHI status between cases ($n = 65$) and controls ($n = 69$) was not statistically significant. In the present study, there were more subjects with good oral hygiene (19 – cases and 17 controls) than subjects with poor oral hygiene (11 – cases and 10 controls) respectively (Table VIII). This may be because there were more females (39 female patients among cases & controls) (Table III). Men have poorer oral hygiene than women, as evidenced by higher levels of plaque & calculus (**U.S. Public Health Service, National Institute of Dental Research 1987**).¹⁰⁸

In the present study *H. pylori* was detected in dental plaque by rapid urease test in patients with good and poor oral hygiene among cases and controls.

H. pylori may be part of normal oral microenvironment and belong to a normal bacterial film that is not a pathogenic reservoir of *H. pylori* for the stomach. However when the host's immunological defense becomes impaired, bacteria's role as commensal is changed and it becomes a pathogen (**M. Czenikiewicz – Guzik et al 2005**).²¹

H.pylori could be a transient microbe which comes in with food and vanishes after some time depending on some host factors as well as its interactions with the residual flora of oral biofilm (**Okuda et al 2000**).⁸⁹

Apart from the oral-oral & faecal-oral route of transmission, another proposed route of transmission of *H. pylori* is gastro – oral (**Axon ATR 1995**).⁶ Gastritis especially the acute stage is often accompanied by increased episodes of intermittent reflux which may facilitate the passage of viable organisms into the mouth (**Axon ATR 1995**).⁶ Given the forcible nature of emesis, plaque biofilm could become impregnated with ejected *H. pylori*,(**Parsonnet 1999**)⁹³ and may serve as a source of infection or reinfection.

The periodontal disease status of the patient was also examined. The diagnosis of chronic periodontitis was based on Armitage classification² 1999. Patients with generalized and localized periodontitis were put together and grouped as those with periodontitis and the remaining were grouped as healthy. Pocket ≥ 5 mm in at least one site was associated with increased odds of *H.pylori* seropositivity in the NHANES Survey (1988-1991). Hence this was also considered as a variable in the present study (Table IX).In the present study there were more subjects with periodontitis among cases than controls. This was found to be statistically significant with a P value of < 0.001 and OR of 2.89 (95% CI 1.06 to 78) by multiple logistic regression analysis (Tables VII and X).

It concurs with the study by **B.A. Dye et al 2002**,²⁹ that poor periodontal health characterized by advanced periodontal pockets may be associated with *H. pylori* infection in adult patients independent of poverty status.

H.pylori survives in moderate to advanced periodontal pockets because the architecture and the microcosm of these periodontal

conditions promote a viable habitat for microaerophilic and anaerobic micro-organisms. Because dental biofilm can provide urea, urease producing bacteria such as *H.pylori* may have improved viability in this periodontal environment.

Correlation analysis was done between oral hygiene status of the patients and presence of *H. pylori* in dental plaque and thus with gastric infection in cases and controls. It was not found to be statistically significant (Table XIII and XIV).The reasons are as given above.

When the periodontal disease status was correlated with oral *H.pylori* and the corresponding gastric infection in cases and controls, it was found that although all the cases of periodontitis were positive for *H.pylori* by RUT, it was not statistically significant ($P=1.00$) in cases but was found to be statistically significant ($P=0.005$) in controls(Table XIII & XIV).In certain oral conditions like periodontitis, mucosal ulcers, the numbers of *H.pylori* may increase and reach levels sufficient to cause infection, especially when the immunological status of the patient is compromised ***Choudary CR et al 2003***.¹⁹Hence the periodontal pocket could serve as a reservoir for future gastric infection in these controls.

The methods of detection of *H.pylori* in dental plaque vary and include the rapid urease test, culture and PCR. In the present study, both the rapid urease test and culture were employed to detect the presence of *H.pylori* in dental plaque. *H. pylori* is the only urease – positive bacteria known to reside in the stomach; hence the rapid urease test (RUT) can be used to confirm the presence of *H. pylori* in gastric samples.

The oral cavity is residence to several urease producing species including *Streptococcus* species, *Haemophilus* species and *Actinomyces* species. But *Vaira D.1988*¹⁰⁹ reported that urease positive micro-organisms present in the oral cavity usually cannot give positive results within an hour. Hence the RUT has been used by several studies to detect the presence of *H.pylori* in dental plaque *Gill et al 1994*³⁹, *Pytko Polonczyk et al 1996*⁹⁶, *Desai et al 1998*²⁵, *Butt et al 1999*¹⁷, *Avcu et al 2001*⁵, *Pradeep et al 2006*.⁹⁴ Studies have reported that the RUT has a specificity near 100% and sensitivity between 70% and 93% (*Mc Nulty 1992*⁷⁵, *Delterse M. 1988*).²⁴ The sensitivity of using the RUT in dental plaque to determine *H.pylori* status is reported to be 89.7% with a diagnostic accuracy of 86.7% (*Gurbuz et al 2003*).⁴³

In the present study, more cases (48 of 55) had positive RUT in the dental plaque than controls (29 of 55) and this was found to be statistically significant with a P value of < 0.001 and OR of 5.05 with 95% CI of 1.71 to 14.89 (by multiple logistic regression analysis) (Tables X and XII). This is similar to the study by *Pradeep et al 2006*,⁹⁴ wherein 58 of 65 cases and 49 of 69 controls were RUT positive in dental plaque. This was found to be statistically significant with an OR of 3.10 (95% CI 1.21 to 9.77).

In the present study, *H.pylori* could not be detected by culture in dental plaque samples of either cases or controls (Table X). This is in accordance with other studies, wherein there was no cultural detection of *H.pylori* in dental plaque samples- *Cheng et al 1996* 0/122¹⁸ dental plaque samples, *Luman et al 1996* 0/109⁵⁹, *Allaker et al 2002* 0/100¹,

Ishihara et al 1997 0/87⁴⁸, *Hardo et al 1995* 0/65⁴⁴ dental plaque samples.

The failure of culture methods to detect H.pylori from dental plaque may be due to the following reasons

- i) Insufficient numbers in oral samples(*Song et al 2000*).¹⁰³
- ii) H.pylori requires a microaerophilic environment, supplemented media and up to 7 days incubation for growth. Under these conditions, overgrowth by other oral species is likely and may inhibit the growth of oral H.pylori (*Ishihara et al 1997*).⁴⁸
- iii) Absence of culturable oral H.pylori and the presence of a viable coccoid form that is unculturable by conventional techniques (*Bode et al 1993*).¹⁶
- iv) Being microaerophilic, these bacteria may not be able to survive the sampling process before being transferred into the medium. This is particularly true, when the numbers of colonizing bacteria are low (*Asikainen S. et al 1994*)⁴.

The results of this study showed that although the rapid urease test was positive for 48 of 55 cases and 29 of 55 controls, culture was negative in the dental plaque samples of all cases and controls. Hence the role of dental plaque as a reservoir for H.pylori infection is inconclusive.

SUMMARY AND CONCLUSION

110 patients with dyspeptic symptoms and clinical indications for an upper gastroendoscopy from the outpatient Department of Medical Gastroenterology Government General Hospital Chennai were selected for the study. The study variables age, gender, handling of animals, smoking and alcohol consumption were obtained from the patient's history. The socio-economic status was arrived, based on Kuppuswamy's Socio-economic status 2007. Oral hygiene status was evaluated using the Simplified Oral Hygiene Index of Greene and Vermillion 1964. Probing depth and clinical attachment level were recorded and based on the AAP 1999 classification, patients were classified into periodontitis and healthy patients. Antral biopsy samples (during endoscopy) and dental plaque samples were obtained. *H.pylori* in dental plaque was detected by RUT and culture, while antral biopsy samples were examined for the presence of *H.pylori* by RUT, culture, histopathology and ELISA. Patients with clinical symptoms and a positive test on antral biopsy sample by any of the three diagnostic tests – RUT, histology, and serology were cases and those negative were controls.

Among the variables, handling of animals was found to be a risk factor for acquiring *H.pylori* infection. The oral hygiene status of the patients did not influence *H.pylori* infection. Patients with periodontitis were at a higher risk for developing *H.pylori* infection and pocket depths $\geq 5\text{mm}$ could serve as a reservoir for the organism. All these observations are based on the positive results obtained with the rapid urease test in

dental plaque. However there was no cultural detection of *H.pylori* in dental plaque samples in this study. Hence the presence of *H.pylori* in dental plaque is inconclusive.

Dental plaque cannot however be discounted as a possible alternate site for the organism. The present methods may be inadequate to reliably isolate the organism from this site. A more comprehensive search for the organism in this environment and other ecological niches within the gingival crevices ought to be conducted to elucidate the role of dental plaque as a potential reservoir for *H.pylori*.

If the oral cavity is a reservoir for gastric infection, even in a minority of individuals, the control of dental plaque along with standard periodontal procedures should be recommended for patients with chronic gastritis or peptic ulcer.

BIBLIOGRAPHY

1. Allaker RP, Young KA, Hardie JM, Domizio P, Meadows NJ (2002). Prevalence of *Helicobacter pylori* at oral and gastrointestinal sites in children: evidence for possible oral-to-oral transmission. *J Med Microbiol* 51:312-317.
2. Armitage GC: Development of a classification system for periodontal diseases and conditions, *Ann Periodontol* 4:1,1999.
3. Ash ASF, Schild HO. Receptors mediating some actions of histamine. *Br J Pharmacol Chemother* 1966; 27: 427-39.
4. Asikainen S, Chen C, Slots J. Absence of *Helicobacter pylori* in subgingival samples determined by PCR. *Oral Microbiol Immunol* 1994; 9: 318-320.
5. Avcu N, Ferit Avcu, Cengiz Beyan et al. the relationship between gastric- oral *Helicobacter pylori* and oral hygiene in patients with vitamin B₁₂ – deficiency anaemia. *Oral Surg Oral Med Oral Path Oral Radiol Endod* 2001; 92: 166-169.
6. Axon ATR: Is *Helicobacter pylori* transmitted by the gastro-oral route? *Aliment Pharmacol Ther* 9: 585- 588, 1995.
7. Bantvala N, Lopez CR, Owen R et al, *Helicobacter pylori* in dental plaque. *Lancet* 1993; 341: 380.
8. Bantvala N, Lopez CR, Owen RJ et al Use of the polymerase chain reaction to detect *Helicobacter pylori* in the dental plaque of healthy and symptomatic individuals. *Microbial Ecol Health Dis* 1994;7:1-8.
9. Berroteran A, Perrone M, Correnti M et al Detection of *Helicobacter pylori* DNA in the oral cavity and gastro duodenal system of a Venezuelan population. *J Med Microbiol* 2002; 51; 764-770.
10. Bielanski W. Epidemiological study on *Helicobacter pylori* infection and extragastrointestinal disorders in Polish population. *J Physiol Pharmacol* 1999; 50:723-733.
11. Bilkoth CAT. Offenes Schreiben an Herrn Dr. L. Wittelshofer. *Wien Med Wochenschr* 1881; 31:161-5, 1427.
12. Birek C, Grandhi R, McNeill K, Singer D, Ficarra G, Bowden G (1999). Detection of *Helicobacter pylori* in oral aphthous ulcers. *J Oral Pathol Med* 28:197-203.
13. Bizzozero G. Sulle ghiandole tubulari del tubo gastroenterico e sui rapporti del loro colPepitelio di rivestimento della mucosa. *Arch Mikr Anat* 1893; 42: 82.

14. Black JW, Duncan WAM, Durant CJ, Ganellin CR, Parsons EM. Definition and antagonism of histamine H₂-receptors. *Nature* 1972; 236: 385-90.
15. Blaser MJ. Ecology of *Helicobacter pylori* in the human stomach. *J Clin Invest* 1997; 100: 759-762.
16. Bode G, Mauch F, Malfertheiner P (1993). The coccoid forms of *Helicobacter pylori*. Criteria for their viability. *Epidemiol Infect* 111: 483-490.
17. Butt AK, Khan AA, Bedi R (1999). *Helicobacter pylori* in dental plaque of Pakistanis. *J Int Acad Periodontol* 78: 78-82.
18. Cheng HH, Webberley M, Evans M, Hanson N, Brown R. *Helicobacter pylori* in dental plaque and gastric mucosa. *Oral Surg Oral Med Oral Pathol Oral Radiol Endol* 81: 421-423.
19. Choudary CR, Choudary AD, Alam S, Markus AF, Tanaka A. Presence of *H. pylori* in the oral cavity of betel – quid (Paan) chewers with dyspepsia: relationship with periodontal health. *Public Health* 2003; 117: 346-347.
20. Coghlan JG, Humphries H, Dooley C et al. *Campylobacter pylori* and recurrence of duodenal ulcers - a 12 month follow-up study. *Lancet* 1987; ii: 109-11.
21. Czesnikiewicz-Guzik.M , Bielanski W., Guzik TJ., Loster B., Konturek SJ., *Helicobacter pylori* in the oral cavity and its implications in gastric infection , periodontal health, immunology and dyspepsia. *J. Physiol Pharmacol* 2005; 56: Suppl 6, 77-89.
22. Czesnikiewicz-Guzik.M ,Karczewska E., Bielanski W., Guzik TJ., Kapera P.et al Association of the presence the *Helicobacter pylori* in the oral cavity and in the stomach. *J. Physiol Pharmacol* 2004; 55: Suppl 2, 105-115.
23. D'Alessandro A, Seri S (1992). Comparison of three different methods for evaluation of *Helicobacter pylori* in human dental plaque. *Boll Soc It Biol Sper* LXVIII: 769-773.
24. Delterse M, BuretteA, Glupcaynski Y. Rapid identification of *Campylobacter pylori* in gastric biopsies. In Menge H, Greysar M, Tygot GNJ, eds. *Campylobacter pylori*. Berlin : Springer Verlag; 1988: 135- 144.
25. Desai HG, Gill HH, Shankaran K, Mehta PR, Prabhu SR (1998). Dental plaque: a permanent reservoir of *Helicobacter pylori*? *Scand J Gastroenterol* 26:1205-1208.
26. Donati M. *De medica htstoria mirabili*. Mantuae per fr. Osanam, lib. IV, Cap iii: 1586; 19.
27. Dowsett S.A., Kowolik M.J. Oral *Helicobacter pylori*: Can We Stomach It? *Crit Rev Oral Biol Med* 2003;14: 5: 226- 233.

28. Dowsett SA, Archila L, Segreto VA, Gonzalez CR, Silva A, Vastola KA, et al (1999). *Helicobacter pylori* infection in indigenous families of Central America: serostatus and oral and fingernail carriage. *J Clin Microbiol* 37:2456-2460.
29. Dye BA, Kruszon- Moran D, McQuillan G. The relationship between periodontal disease attributes and *Helicobacter pylori* infection among adults in the United States. *Am J Public Health* 2002; 92: 1809-1815.
30. Editorial. *Campylobacter pylori* becomes *Helicobacter pylori*. *Lancet* 1989; ii: 1019-2.
31. European *Helicobacter pylori* Study Group 1997. Current European concepts in the management of *Helicobacter pylori* infection. The Maastricht consensus report. *Gut* 41:8- 13.
32. Everhart JE, Kruszon- Moran D, Perez-Perez GI, Tralka T, Mc Quillan G. Seroprevalence and ethnic differences in *Helicobacter pylori* infection among adults in the United States. *J Infect Dis.* 2000; 181: 1359- 1363.
33. Fitzgerald D, Murphy P. Studies in the physiological chemistry and clinical significance of urease and urea with special reference to the stomach. *IT J Med Sci* 1950; 292: 97-15.
34. Fox JG. Non-human reservoirs of *Helicobacter pylori* *Aliment Pharmacol Ther* 1995; 9 Suppl. 2: 93-103.
35. Francis Megraud, Philippe Lehours *Helicobacter pylori* Detection and Antimicrobial Susceptibility Testing. *Clinical Microbiology Reviews* 2007; 20,2: 280-322.
36. Gill HH, Shankaran K, Desai HG. *Helicobacter pylori* in dental plaque of children and their family members. *J Assoc Physicians India* 1994; 42: 719-721.
37. Ganser AL. Forte JG. K'-stimulated ATPase in purified microsomes of bullfrog oxyntic cells. *Biochim Biophys Acta* 1973; 307: 169-80.
38. Gasbarrini G, Pretolani S, Bonvinci F et al. A population based study of *Helicobacter pylori* infection in a European country: the San Marino study. Relations with gastrointestinal diseases. *Gut* 1995; 36:838-844.
39. Gill HH, Shankaran K, Desai HG (1994). *Helicobacter pylori* in dental plaque of children and their family members. *J Assoc Physicians India* 42:719-721.
40. Goodwin CS, Armstrong JA, Chilvers T et al. Transfer of *Campylobacter pylori* and *Campylobacter mustelae* to *Helicobacter* gen. nov. as *Helicobacter pylori* comb. nov. and *Helicobacter mustelae* comb. nov., respectively. *Int J System Bacteriol* 1989; 39: 397-405.

41. Grant DA, Irving B, Stern, Listgarten MA, *Periodontics* 6th edition 525-572.
42. Greene JC, Vermillion JR. The simplified oral hygiene index. *J Am Dent Assoc* 1964; 68: 7-13.
43. Gurbuz AK, Ozel AM, Yazgan Y, Celik M, Yildirim S. Oral Colonization of *Helicobacter pylori*: Risk factors and response to eradication therapy. *South Med J*. 2003; 96: 244-247.
44. Hardo PG, Tugnait A, Hassan F, Lynch DAF, West AP, Mapstone NP, et al (1995). *Helicobacter pylori* infection and dental care. *Gut* 37:544-546.
45. Haruma K, Okamoto S, Kawaguchi H et al : Reduced incidence of *Helicobacter pylori* infection in young Japanese persons between the 1970s and the 1990s . *J Clin Gastroenterol* 25: 583- 586, 1997.
46. International Agency for Research on Cancer. Schistosomes, liver flukes and *Helicobacter pylori*. *IARC Monographs on the Evaluation of Carcinogenic Risk to Humans, Vol 6. Lyon:IARC, 1994.*
47. Isabelle.M. Madinier, Thierry M. Fosse and Roger A. Monteil Oral Carriage of *Helicobacter pylori*: A Review *J. Periodontol* 1997; 68: 2-6.
48. Ishihara K, Miura T, Kimizuko R, Ebihara Y, Mizumo Y, Okuda K(1997). Oral bacteria inhibit *Helicobacter pylori* growth. *FEMS Microbiol Lett* 152: 355-361.
49. Jones DM, AM Lessels and J. Eldridge 1984. *Campylobacter* like organisms on the gastric mucosa: culture, histological and serological studies. *J. Clin Pathol.*37: 1002- 1006.
50. Khandaker MAK, Scott A, Eastwood MA, Palmer KR. Do teeth predispose to duodenal ulcer relapse? *Gut* 1991; 32: A1207.
51. Kim N, Lim SH, Lee KH, You JY, Kim JM, Lee NR, et al (2000). *Helicobacter pylori* in dental plaque and saliva *Korean J Int Med* 15: 187- 194.
52. Krajden S, Fuksa M, Anderson J, Kempston J, Boccia A, Petrea C, et al (1989) Examination of human stomach biopsies, saliva, and dental plaque for *Campylobacter pylori*. *J Clin Microbiol* 27:1397-1398.
53. Krieiutz U. Ueber das Auftreten von Spirochaeten verschiedener Form im Mageninhalt bei *Carcinoma ventricuh*. *Dtsch Med Wochenschr* 1906; 32: 872.
54. Kuipers E, A. Pena, G. Van Kamp et al 1993. Seroconversion for *Helicobacter pylori*. *Lancet* 342: 328-331.

55. Kumar N, C. Shekar, P. Kumar and A.S. Kundu. Kuppasamy, s Socioeconomic Status- Updating for 2007. *The Indian Journal of Pediatrics* 2007; 74: 12: 1131-1132.
56. Lee J.M., Breslin N.P., Fallon C, O' Morain CA. Rapid urease test lacks sensitivity in *Helicobacter* diagnosis when peptic ulcer disease presents with bleeding. *American Journal of Gastroenterology* 2000; 95(5): 1166-1170.
57. Li C, Musich PR, Ha T, et al High prevalence of *Helicobacter pylori* in saliva demonstrated by a novel PCR assay. *J Clin Pathol* 1995; 48: 662-668
58. Luck JM, Seth TN. The physiology of gastric urease. *Biochem J* 1924; 18: 357-65.
59. Luman W, Alkout AM, Blackwell CC, Weir DM, Palmer KR (1996). *Helicobacter pylori* in the mouth—negative isolation from dental plaque and saliva. *Eur J Gastroenterol Hepatol* 8: 11-14.
60. Mackei & Mc'Cartney, *Practical Medical Microbiology* 14th edn.
61. Mai U, Borsch G, Opferkuch W. Route of infection in *Campylobacter pylori* (CP) disease. *Gastroenterology* 1989; 96(suppl 5): 315.
62. Majmudar P, Shah SM, Dhunjibhoy KR, Desai HG (1990). Isolation of *Helicobacter pylori* from dental plaques in healthy volunteers. *Indian J Gastroenterol* 9: 271-272.
63. Malathy HM , Evans Jr DJ, Abramovitch K, Evans DG, Graham DY. *Helicobacter pylori* infection in dental workers: a seroepidemiology study. *Am J Gastroenterol* 1992; 87: 1728-1731.
64. Malfertheiner P, Megraud F, O'Morain C *et al.* Current European concepts in the management of *H. pylori* infection - The Maastricht Consensus Report. *Eur J Gastroenterol Hepatol* 1997; 9: 1-2.
65. Mapstone NP, Lynch D et al The polymerase chain reaction in the diagnosis of *Helicobacter* infection. *Ital J gastroenterol* 1991; 23 suppl 4.
66. Mapstone NP, Lynch DA et al Identification of *Helicobacter pylori* DNA in the mouths and stomachs of patients with gastritis using PCR. *J Clin Pathol* 1993; 46: 540-543.
67. Marianne Quiding- Jarbrink, Mogens Bove & Gunnar Dahlen Infections of the esophagus and the stomach. *Periodontology* 2009; 49: 166-178.
68. Marsh PD 2004. Dental Plaque as a Microbial Biofilm. *Caries Res* 2004; 38: 204.

69. Marshall BJ, Armstrong JA, McGeachie DB, Glancy RJ. Attempt to fulfill Koch's postulates for pyloric *Campylobacter*. *Med J Aust* 1985; 142: 436-9.
70. Marshall BJ, Warren JR. Unidentified curved bacilli in the stomach of patients with gastritis and peptic ulceration. *Lancet* 1984; 1: 1311-1315.
71. Marshall FJ, Goodwin CS. Revised nomenclature of *Campylobacter pyloridis*. *Int J System Bacteriol* 1987; 37: 68.
72. Marshall FJ, Royce H, Annear DL et al. Original isolation of *Campylobacter pyloridis* from human gastric mucosa. *Microbiol Lett* 1984; 25: 83-88.
73. Martin JM Buckley and Colm A O' Morain *Helicobacter* biology-discovery *British Medical Bulletin* 1998; 54, 1:7- 16.
74. Mayo Robson AW, Moynihan BGA. Diseases of the stomach and their surgical treatment. 2nd edn. London: Bailiere, Tindall and Cox, 1904: 165.
75. McNulty CAM. Detection of *Helicobacter pylori* by the biopsy urease test. In: Rathbone BJ, Heatley RV eds *Helicobacter pylori* and *Gastro duodenal Diseases* Oxford: Blackwell Scientific; 1992:58-59.
76. Mealey BL 1999: Influence of periodontal infections on systemic health, *Periodontol* 2000; 21: 197, 1999.
77. Megraud F, Bonnet F, Garnier M, Lamouliatte H. Characterization of *Campylobacter pyloridis* by culture, enzymatic profile and protein contents. *J. Clin Microbiol* 1985; 22: 1007-1010.
78. Mendall MA, Goggin PM, Molineaux N, Levy J, Toosy T, Strachan D, et al. (1992). Childhood living conditions and *Helicobacter pylori* seropositivity in adult life. *Lancet* 339:896-897.
79. Michael JG Farthing *Helicobacter pylori* infection: an overview *British Medical Bulletin* 1998; 54(No.1):1-6.
80. Mitchell HM, Bohane TD et al (1987). Antibody to *Campylobacter pylori* in families of index children with gastrointestinal illness due to *C. pylori*. *Lancet* 681- 682.
81. Mobley H.L.T., M.J. Cortesia, L.E. Rosenthal, and B.D. Jones. Characterization of urease from *Campylobacter pylori*. *J. Clin. Microbiol.* 1988; 26: 831-836.
82. Morgagni GB. *De sedibus et causis morborum per anatomen indagatis, libri quinque*. Venetiis, typog. Remondiniana, 2 vols, 1761.

83. Mravak-Stipetic M, Gall-Troselj K, Lukac J, Kusic Z, Pavelic K, Pavelik J (1998). Detection of *Helicobacter pylori* in various oral lesions by nested polymerase chain reaction (PCR). *J Oral Pathol Med* 27:1-3.
84. Nabwera HM, Logan RP. Epidemiology of *Helicobacter pylori*: transmission and extragastric reservoirs. *J Physiol Pharmacol* 1999; 50: 711- 722.
85. Namavar F, Roosendaal R, Kuipers EJ, de Groot P, van der Bijl MW, Pena AS, et al. (2001). Presence of *Helicobacter pylori* in the oral cavity, oesophagus, stomach and faeces of patients with gastritis. *Eur J Clin Microbiol Infect Dis* 14:234.
86. Newman MG, Takei HH, Klokkevold PR, Carranza FA, *Clinical Periodontology* 10th edition 552-553.
87. Newman MG, Takei HH, Klokkevold PR, Carranza FA, *Clinical Periodontology* 10th edition 494.
88. NIH Consensus Development Panel on *Helicobacter pylori* in peptic ulcer disease. *Helicobacter pylori* in peptic ulcer disease. *JAMA* 1994; 272: 65-9.
89. Okuda K, Ishihara K, Miura T et al *Helicobacter pylori* may have only a transient presence in the oral cavity and on the surface of oral cancer. *Microbiol Immunol* 2000; 44: 385-388.
90. Oshowo A, Tunio M, Gillam D, Botha AJ, Holton J, Boulos P, et al. (1998). Oral colonization is unlikely to play an important role in *Helicobacter pylori* infection. *Br J Surg* 85:850-852.
91. Ozdemir A., Mas M.R., Sahin S., Saglamkaya U., Atekesan U. Detection of *Helicobacter pylori* colonization in dental plaques and tongue scrapings of patients with chronic gastritis. *Quintessence Int* 2001; 32:131- 134.
92. Palmer ED. Investigation of the gastric spirochetes of the human. *Gastroenterology* 1954; 27:218-20.
93. Parsonnet J, Shmueli H, Haggerty BS (1999). Fecal and oral shedding of *Helicobacter pylori* from healthy infected adults. *J Am Med Assoc* 282:2240-2245.
94. Pradeep S. Anand, K. Nandakumar, and K.T. Shenoy. Are Dental Plaque, Poor Oral Hygiene, and Periodontal Disease Associated With *Helicobacter pylori* Infection? *J Periodontal* 2006; 77, 4: 692- 698.
95. Pustorino R, Nicosia R, D'Ambra G, Di Paola M, Brugnoletti O, Grippaudo G, et al. (1996). The mouth-stomach crossing of *Helicobacter pylori*. *Eur Rev Med Pharmacol Sci* 18:183-186.

96. Pytko-Polonczyk J, Konturek SJ, Karczewska E, Bielanski W, Kaczmarczyk-Stachowska A (1996). Oral cavity as a permanent reservoir of *Helicobacter pylori* and potential source of reinfection. *J Physiol Pharmacol* 47:121-129.
97. R.Nijhawan, R.Kochhar, D.Panigrahi et al. Identification of *Helicobacter pylori* by Endoscopic Crush Cytology. *Indian J. Gastroenterol* 1993; 12(2): 45-46.
98. Riggio MP, Lennon A. Identification by PCR of *Helicobacter pylori* in subgingival plaque of adult periodontitis. *J Med Microbiol.* 1999 ;48 : 317- 322.
99. Roger .A. Feldman, A James P Eccersley and Jeremy M. Hardie Epidemiology of *Helicobacter pylori*: acquisition, transmission, population prevalence and disease to- infection ratio. *British Medical Bulletin* 1998; 54, 1: 39- 53.
100. Schwarz K. Ueber penetrierende Magen- und Jejunalgeschwure. *Beitr Klin Chir* 1910; 67:96-128.
101. Sippy BW. Gastric and duodenal ulcer. Medical cure by an efficient removal of gastric juice *JAMA* 1915; 64: 1625-30.
102. Song Q, Haller B, Ulrich D, Wichelhaus A, Adler G, Bode G (2000). Quantification of *Helicobacter pylori* in dental plaque samples by competitive polymerase chain reaction. *J Clin Pathol* 53: 218-222.
103. Steer HW. Infrastructure of cell migration through the gastric epithelium and its relationship to bacteria. *J Clin Pathol* 1975; 28: 639-46.
104. Tannock GW. Normal microflora. An introduction to microbes inhabiting the human body. London: Chapman and Hall, 1995.
105. Taylor DN, Blaser MJ (1991). The epidemiology of *Helicobacter pylori* infection. *Epidemiol Rev* 13:42-59.
106. The EUROGAST Study Group. Epidemiology of, and risk factors for, *Helicobacter pylori* infection among 3194 asymptomatic patients in 17 populations. *Gut* 1993; 34: 1672-1676.
107. Unge P. Review of *Helicobacter pylori* eradication regimens. *Scand J Gastroenterol Suppl* 1996; 215: 74-81.
108. U.S. Public Health Service, National Centre for Health Statistics: *Oral hygiene in adults, United States, 1960-1962*, PHS Pub No 1000, Series 11, No 16, Washington, DC, 1966, US Government Printing Office.
109. Vaira D, Holton J, Cairns et al. Urease test for *Campylobacter pylori*: Care in interpretation. *J Clin Pathol* 1988; 41: 812- 813.

110. Veldhuyzen van Zanten SJ. Do socioeconomic status, marital status and occupation influence the prevalence of *Helicobacter pylori* infection? *Aliment Pharmacol Ther* 1995; 9 Suppl. 2 : 41-44.
111. Von Mikulicz-Radecki J. Ueber Gastroskopie und Aesophagoskopie. *Wien Med Press* 1881; 22:1405-8, 1437[^]3, 1473-5, 1505-7, 1537-41, 1573-7, 1629-31.
112. Warren JR, Marshall B. Unidentified curved bacilli on gastric epithelium in active gastritis. *Lancet* 1983; i: 1273-1275.
113. Yang HT. Nested polymerase chain reaction in detection of *Helicobacter pylori* in human dental plaque. *Chin Med J* 1993; 73:750-752.

APPENDIX PROFORMA

ASSOCIATION OF DENTAL PLAQUE, POOR ORAL HYGIENE AND PERIODONTAL DISEASE WITH HELICOBACTER PYLORI INFECTION.

Date:

O.P.No:

Name:

Code No:

Age/Sex:

Address:

Tel.No:

Occupation:

Income:

Level of education:

SES score:

Chief complaints:

Duration:

History:

Smoker – Current/Ex-smoker/ Not at all

Alcohol– Current/Past/ Never

H/o contact with animals- Yes/No

Type of animals-

SIMPLIFIED ORAL HYGIENE INDEX- GREENE AND VERMILION 1964

DEBRIS INDEX

16	11	26
46	31	36

$$DI = \frac{\text{Sum of debris score per tooth surface}}{\text{No. of surfaces examined}}$$

CALCULUS INDEX

16	11	26
46	31	36

$$CI = \frac{\text{Sum of calculus score per tooth surface}}{\text{No. of surfaces examined}}$$

OHI score= Debris index score+ Calculus index score
Interpretation- Good/Fair/ Poor oral hygiene

PROBING DEPTH (PPD) AND CLINICAL ATTACHMENT LOSS (CAL) (in mm) (CARRANZA)

MAXILLARY **Buccal**

[illegible]

Palatal

MANDIBULAR **Buccal**

[illegible]

Lingual

Periodontal disease status – H- Healthy G- Generalized chronic periodontitis L- Localized chronic periodontitis

RUT (P):

RUT (B):

CULTURE (P):

CULTURE (B):

GIEMSA STAIN:

GRAM STAIN:

ELISA:

HISTOPATHOLOGICAL FINDINGS:

INFORMED CONSENT FORM-ENGLISH

Study Title:

ASSOCIATION OF DENTAL PLAQUE, POOR ORAL HYGIENE AND PERIODONTAL DISEASE WITH HELICOBACTER PYLORI INFECTION

Date:

O.P. No:

Name:

Code No:

Address:

Age/Sex:

Tel. No:

I, _____ age _____ yrs, exercising my free power of choice, hereby give my consent to be included as a participant in the clinical study “Association of dental plaque, poor oral hygiene and periodontal disease with Helicobacter pylori infection”.

I agree to the following:

I have been informed to my satisfaction about the purpose of the study, and nature of the study.

I understand that I should undergo endoscopy procedure as a part of diagnosis for my condition in the stomach. The tissue sample taken during the procedure for the purpose of diagnosis will be used for study purpose and helps in confirming the diagnosis.

I understand that the investigations will also require oral plaque sample in required amounts.

I agree to co-operate fully and participate in the study.

I hereby give permission to use the records for study. I am told that study doctor and study institution will keep my identity confidential.

Signature of the
Investigator

Signature of the
Guide

Signature of the
Participant

ஆராய்ச்சி ஒப்புதல் படிவம்

ஆராய்ச்சி தலைப்பு:

”எச்.பைலோரி நுண்ணுயிர் தொற்றுள்ள நிலையில் பற்புறத்திசு நோய்க்கும், பற்கிருமி படலம் மற்றும் பராமரிக்கப்படாத வாய்சுகாதாரம் குறித்து தொடர்பு”.

தேதி : புறநோயாளி எண் :
பெயர் : ஆராய்ச்சி சேர்க்கை எண் :
முகவரி : வயது : ஆ/பெ:
தொலைபேசி எண் :

நான் _____ வயது _____ என்னுடைய சுயநினைவுடன் மற்றும் முழு
சுதந்திரத்துடன் இந்த மருத்துவ ஆராய்ச்சியில் என்னை சேர்த்துக் கொள்ள
சம்மதிக்கிறேன்.

எனக்கு விளக்கப்பட்ட விஷயங்களுக்கு நான் எனது சம்மதத்தை தருகிறேன்

- இந்த ஆராய்ச்சியின் நோக்கம் மருத்துவ முறைகள் பரிசோதனை முறைகள் எனக்கு திருப்தியுறும் வகையில் விளக்கப்பட்டன.
- நோய் அறியும் செயலின் ஒரு பகுதியாக எண்டோஸ்கோபி சோதனைக்கு நான் உட்பட வேண்டும் என்பதை உணர்ந்து கொண்டேன். மேலும் மேற்கண்ட சோதனையின் திசுப்பரிசோதனைக்காக சிறிய அளவில் திசு எடுக்கப்படும் எனவும் அது நோயை அறிய உதவும் எனவும், எந்த விதத்திலும் அது சிகிச்சையை பாதிக்காது எனவும் உணர்ந்து கொண்டேன். மேலும் வழக்கமாக செய்யும் இரத்தப் பரிசோதனையும் இந்த ஆய்வுக்கு உபயோகப்படுத்தப்படுகிறது.
- கிருமியை பரிசோதனை செய்வதற்காக பற்களில் படிந்திருக்கும் பற்படலத்தில் சிறிதளவு எடுக்க வேண்டியுள்ளதாக அறிகிறேன்.
- எனது மருத்துவ குறிப்பேடுகளை இந்த ஆராய்ச்சியில் பயன்படுத்திக் கொள்ள சம்மதிக்கிறேன். ஆராய்ச்சி மையமும் ஆராய்ச்சியாளரும் என்னுடைய பெயர் மற்றும் சில விபரங்களை இரகசியமாக வைப்பதாக அறிகிறேன்.

பேராசியரின்
கையொப்பம்

ஆராய்ச்சியாளரின்
கையொப்பம்

நோயாளின்
கையொப்பம்/கைநாட்டு